

Notation: $B = \{z \in \mathbb{C} : |z| < 1\}$, the open unit disc

1. Let D be a connected open set in \mathbb{C} and f a function analytic on D .
 - a) Assuming that f has an analytic logarithm in D , show that f has analytic m th roots for all m . Explicitly: Assume that there exists h analytic on D so that $f = e^h$, and show that for each $m \in \mathbb{N}$ there exists a function g_m analytic on D so that $f = (g_m)^m$.
 - b) Assume now that $f \neq 0$ and f has analytic m th roots for all m . Show that $f(z) \neq 0$ for all $z \in D$. Hint: Look at multiplicities.
2. (a) Prove that if a, b, c and d are real numbers satisfying $ad - bc = 1$, then

$$M(z) = \frac{az + b}{cz + d}$$

maps the upper half-plane $\mathbb{H} = \{x + iy : y > 0\}$ conformally onto itself.

(b) Let $f: B \rightarrow \mathbb{C}$ be such that the imaginary part of f is non-negative for all z in B . Assume that f is non-constant and is analytic in B . Show that

- (i) the imaginary part of f is strictly positive for all z in B ;
- (ii) if $f(0) = i$, then

$$|f(z)| \leq \frac{1 + |z|}{1 - |z|}.$$

Hint: Use the Schwarz Lemma.

3. (a) Suppose f is an entire function such that f^2 is a polynomial. Does it follow that f is a polynomial?
 - (b) Suppose f is analytic on B and f^2 is a polynomial on B . Does it follow that f is a polynomial on B ?
- (Of course, you have to explain your answers).

4. Use residue theory to calculate with justification that

$$\int_0^\infty \frac{x \sin x}{x^2 + 4} dx = \frac{\pi}{2e^2}.$$

5. (a) Let G denote an open set in the complex plane containing \bar{B} . Suppose f is analytic on G and is non-zero on ∂B . Show that, with the counterclockwise orientation for ∂B ,

$$\frac{1}{2\pi i} \int_{\partial B} \frac{f'(z)}{f(z)} dz = N(f, B),$$

where $N(f, B)$ counts the number of zeros of f in B with multiplicity.

(b) Suppose f_n is a sequence of functions analytic on B such that $f_n \rightarrow f$ uniformly on B . Suppose f has exactly R zeros in the set $D = \{z : |z| < \frac{1}{2}\}$ and no zeros on ∂D . Show that there exists an $N > 0$ such that, for all $n > N$, the function f_n has exactly R zeros in D .

6. By definition, an *automorphism* of a connected open set G is a one-to-one analytic function mapping G onto G .

(a) Let f and g be automorphisms of B . Suppose there are two distinct points P and Q in B such that $f(P) = g(P)$ and $f(Q) = g(Q)$. Show that $f = g$ on B .

(b) Let D be a simply connected domain in \mathbb{C} which is not equal to \mathbb{C} , and let f and g be automorphisms of D . Suppose there are two distinct points P and Q in D such that $f(P) = g(P)$ and $f(Q) = g(Q)$. Show that $f = g$ on D .