

Reduction of Order

Recall that the general solution of a second order homogeneous linear differential equation

$$(15.1) \quad L[y] = y'' + p(x)y' + q(x)y = 0$$

is given by

$$(15.2) \quad y(x) = c_1 y_1(x) + c_2 y_2(x)$$

where y_1 and y_2 are any two solutions such that

$$(15.3) \quad W[y_1, y_2](x) = y_1(x)y_2'(x) - y_1'(x)y_2(x) \neq 0 \quad .$$

In this section we shall assume that we have already found one solution y_1 of (15.1) and that we are seeking to find another solution y_2 so that we can write down the general solution as in (15.2).

So suppose we have one non-trivial solution $y_1(x)$ of (15.1) and suppose there is another solution of the form

$$(15.4) \quad y_2(x) = v(x)y_1(x) \quad .$$

Then

$$(15.5) \quad \begin{aligned} W[y_1, y_2] &= y_1 y_2' - y_1' y_2 \\ &= y_1 (v' y_1 + v y_1') - y_1' (v y_1) \\ &= (y_1)^2 v' \\ &\neq 0 \end{aligned}$$

unless $v' = 0$. Thus, any solution we construct by multiplying our given solution $y_1(x)$ by a non-constant function $v(x)$ will give us another linearly independent solution.

The question we now wish to address is: how does one find an appropriate function $v(x)$?

Certainly, we want to choose $v(x)$ so that $y_2(x) = v(x)y_1(x)$ satisfies (15.1). So let us insert $y(x) = v(x)y_1(x)$ into (15.1):

$$(15.6) \quad \begin{aligned} 0 &= \frac{d^2}{dx^2} (v y_1) + p(x) \frac{d}{dx} (v y_1) + q (v y_1) \\ &= v'' y_1 + 2v' y_1' + v y_1'' + p(x) v' y_1 + p(x) v y_1' + q v y_1 \\ &= v (y_1'' + p(x) y_1' + q(x) y_1) + v'' y_1 + (2y_1' + p(x) y_1) v' \end{aligned}$$

The first term vanishes since y_1 is a solution of (15.1), so $v(x)$ must satisfy

$$(15.7) \quad 0 = y_1 v'' + (2y_1' + p(x) y_1) v'$$

or

$$(15.8) \quad v'' + \left(p(x) + \frac{2y_1'}{y_1} \right) v' = 0 \quad .$$

Now set

$$(15.9) \quad u(x) = v'(x) \quad .$$

Then we have

$$(15.10) \quad u' + \left(p(x) + \frac{2y_1'(x)}{y_1(x)} \right) u = 0 \quad .$$

This is a first order linear differential equation which we know how to solve. Its general solution is

$$(15.11) \quad \begin{aligned} u(x) &= C \exp \left[- \int^x \left(p(t) + \frac{2y_1'(t)}{y_1(t)} \right) dt \right] \\ &= C \exp \left[- \int^x p(t) dt - 2 \int^x \frac{y_1'(t)}{y_1(t)} dt \right] \end{aligned}$$

Now note that

$$(15.12) \quad \frac{d}{dt} \ln [y_1(t)] = \frac{y_1'(t)}{y_1(t)} \quad ,$$

so

$$(15.13) \quad \begin{aligned} \exp \left[-2 \int^x \frac{y_1'(t)}{y_1(t)} dy \right] &= \exp \left[-2 \int^x \frac{d}{dt} (\ln [y_1(t)]) dt \right] \\ &= \exp \left[-2 \ln [y_1(x)] \right] \\ &= \exp \left[\ln [(y_1(x))^{-2}] \right] \\ &= \frac{1}{(y_1(x))^2} \quad . \end{aligned}$$

Thus, (15.11) can be written as

$$(15.14) \quad u(x) = \frac{C}{(y_1(x))^2} \exp \left[- \int^x p(t) dt \right] \quad .$$

Now recall from (15.9) that $u(x)$ is the derivative of the factor $v(x)$ which we originally sought out to find. So

$$(15.15) \quad \begin{aligned} v(x) &= \int^x u(t) dt + D \\ &= \int^x \left[\frac{C}{(y_1(t))^2} \exp \left[- \int^t p(t') dt' \right] \right] + D \end{aligned}$$

It is not too difficult to convince oneself that it is not really necessary to carry along the constants of integration C and D . For the constant D can be absorbed into the constant c_1 of the general solution $y(x) = c_1 y_1(x) + c_2 y_2(x)$, while the factor C can be absorbed into the constant c_2 . Thus, without loss of generality, we can take $C = 1$ and $D = 0$. So given one solution $y_1(x)$ of (15.1), a second solution $y_2(x)$ of (15.1) can be formed by computing

$$(15.16) \quad v(x) = \int^x u(t) dt$$

where

$$(15.17) \quad u(x) = \frac{1}{(y_1(x))^2} \exp \left[- \int^x p(t) dt \right]$$

and then setting

$$(15.18) \quad y_2(x) = v(x) y_1(x) \quad .$$

The general solution of (15.1) is then

$$(15.19) \quad y(x) = c_1 y_1(x) + c_2 v(x) y_1(x) \quad .$$

This technique for constructing the general solution from single solution of a second order linear homogeneous differential equation is called **reduction of order**.

For those of you who like nice tidy formulae we can write

$$(15.20) \quad \boxed{y_2(x) = y_1(x) \int^x \frac{1}{(y_1(s))^2} \exp \left[- \int^s p(t) dt \right] ds}$$

for the second solution.

EXAMPLE 15.1.

$$(15.21) \quad y_1(x) = e^{-x}$$

is one solution of

$$(15.22) \quad y'' + 2y' + y = 0 \quad .$$

Find another linearly independent solution.

Well, $p(x) = 2$, so

$$(15.23) \quad \begin{aligned} u(x) &= \frac{C}{(y_1(x))^2} \exp \left[- \int^x p(t) dt \right] \\ &= \frac{C}{e^{-2x}} \exp[-2x] \\ &= C e^{2x} e^{-2x} \\ &= C \end{aligned}$$

So

$$(15.24) \quad v(x) = \int^x u(t) dt = \int^x C dt = Cx.$$

Thus,

$$(15.25) \quad y_2(x) = v(x)y_1(x) = Cxe^{-x} \quad .$$

EXAMPLE 15.2.

$$(15.26) \quad y_1(x) = x$$

is a solution of

$$(15.27) \quad x^2 y'' + 2xy' - 2y = 0 \quad .$$

Use reduction of order to find the general solution.

Well, we first put the differential equation in standard form:

$$(15.28) \quad y'' + \frac{2}{x}y' - \frac{2}{x^2}y = 0 \quad .$$

Thus,

$$(15.29) \quad p(x) = \frac{2}{x}, \quad q(x) = \frac{-2}{x^2} \quad .$$

We first compute $u(x)$.

$$(15.30) \quad \begin{aligned} u(x) &= \frac{C}{(y_1(x))^2} \exp \left[- \int^x p(t) dt \right] \\ &= \frac{C}{x^2} \exp \left[\int^x -\frac{2}{t} dt \right] \\ &= \frac{C}{x^2} \exp[-2 \ln|x|] \\ &= \frac{C}{x^2} x^{-2} \\ &= Cx^{-4} \quad . \end{aligned}$$

So

$$(15.31) \quad \begin{aligned} v(x) &= \int^x u(t) dt \\ &= \int^x Ct^{-4} dt \\ &= -\frac{C}{3} x^{-3} \end{aligned}$$

and

$$(15.32) \quad y_2(x) = v(x)y_1(x) = -\frac{C}{3x^3}x = C'x^{-2} \quad .$$

The general solution of the original differential equation is thus

$$(15.33) \quad y(x) = c_1 y_1(x) + c_2 y_2(x) = c_1 x + c_2 x^{-2} \quad .$$