

The Laplace Transform

Suppose $f : \mathbb{R} \rightarrow \mathbb{R}$ is a “nice” (to be qualified latter) function of x . The **Laplace transform** $\mathcal{L}[f]$ of f is the function from \mathbb{R} to \mathbb{R} defined by

$$(28.1) \quad \mathcal{L}[f](s) = \int_0^{\infty} e^{-sx} f(x) dx \quad .$$

We note that in the formula above, s is the variable upon which the Laplace transform $\mathcal{L}[f]$ depends.

EXAMPLE 28.1. If

$$(28.2) \quad f(x) = ax$$

then

$$(28.3) \quad \begin{aligned} \mathcal{L}[f](s) &= \int_0^{\infty} axe^{-sx} dx \\ &= \lim_{N \rightarrow \infty} \left(-\frac{a}{s} xe^{-sx} - \frac{a}{s^2} e^{-sx} \right) \Big|_0^N \\ &= \frac{a}{s^2} \end{aligned}$$

Note that this result really only makes sense for $s > 0$; for $x \leq 0$ the integral does not converge.

EXAMPLE 28.2. If

$$(28.4) \quad f(x) = \sin(ax)$$

then, integrating by twice by parts,

$$(28.5) \quad \begin{aligned} \mathcal{L}[f](s) &= \int_0^{\infty} \sin(ax)e^{-sx} dx \\ &= \lim_{N \rightarrow \infty} \left(e^{-sx} \frac{1}{a} \cos(ax) \right) \Big|_0^N + \frac{s}{a} \int_0^{\infty} e^{-sx} \cos(ax) dx \\ &= \frac{1}{a} + \frac{s}{a} \int_0^{\infty} e^{-sx} \cos(ax) dx \\ &= \frac{1}{a} + \lim_{N \rightarrow \infty} \left(-\frac{1}{a} e^{-sx} \sin(ax) \right) \Big|_0^N - \frac{s^2}{a^2} \int_0^{\infty} e^{-sx} \sin(ax) dx \\ &= \frac{1}{a} + 0 - \frac{s^2}{a^2} \mathcal{L}[f](s) \quad , \end{aligned}$$

we find

$$(28.6) \quad \mathcal{L}[f](s) = \frac{a}{1 + \frac{s^2}{a^2}} = \frac{a}{a^2 + s^2} \quad .$$

(If $s \leq 0$, the integral on the first line does not converge, so $\mathcal{L}[f](s)$ is only defined for $s > 0$.)

EXAMPLE 28.3. If $f(x) = e^{bx}$, then

$$(28.7) \quad \begin{aligned} \mathcal{L}[f] &= \int_0^{\infty} e^{bt} e^{-st} dt \\ &= \int_0^{\infty} e^{(b-s)t} dt \\ &= \frac{1}{b-s} e^{(b-s)t} \Big|_0^{\infty} \\ &= \frac{1}{s-b} \quad (\text{if } s > b) \end{aligned}$$

(If $s \leq b$ then the integral does not converge.)

The following theorem explains under what conditions we can expect the Laplace transform of a function $f(x)$ to exist.

THEOREM 28.4. Suppose that $f(x)$ is a piecewise continuous function for $0 \leq t \leq A$ and there exist constants K, a, M such that

$$(28.8) \quad |f(t)| \leq Ke^{at} \quad , \quad \forall t > M > 0 \quad .$$

Then the Laplace transform $\mathcal{L}[f]$ defined by

$$(28.9) \quad \mathcal{L}[f](s) = \int_0^{\infty} f(t)e^{-st} dt$$

exists for all $s > a$.

The condition (28.8) is a rather moderate “growth” condition on the function $f(x)$; it says that for large enough t , $|f(t)|$ grows no faster than an exponential function of the form Ke^{at} . This condition is easily satisfied by any polynomial function of x .

THEOREM 28.5. **Properties of the Laplace Transform**

(i) Suppose $f_1(x)$ and $f_2(x)$ are two functions satisfying the hypotheses of Theorem 6.2. Then if $g(x) = c_1f_1(x) + c_2f_2(x)$, $\mathcal{L}[g]$ exists and

$$(28.10) \quad \mathcal{L}[g](s) = c_1\mathcal{L}[f_1](s) + c_2\mathcal{L}[f_2](s) \quad .$$

(ii) Suppose that f is continuous and that both f and its derivative f' satisfy the hypotheses of Theorem 6.2. Then $\mathcal{L}[f'](s)$ exists for $s > a$ and moreover

$$(28.11) \quad \mathcal{L}[f'] = s\mathcal{L}[f] - f(0) \quad .$$

(iii) Suppose that f and its derivatives $f', \dots, f^{(n-1)}$ are continuous and satisfy the hypotheses of Theorem 6.2. Then $\mathcal{L}[f^{(n)}](s)$ exists for $s > a$ and

$$(28.12) \quad \mathcal{L}[f^{(n)}](s) = s^n\mathcal{L}[f](s) - s^{n-1}f(0) - s^{n-2}f'(0) - \dots - sf^{(n-2)}(0) - f^{(n-1)}(0) \quad .$$

Proof of (i).

This follows from the linearity property integration:

$$(28.13) \quad \begin{aligned} \mathcal{L}[c_1f_1 + c_2f_2](s) &= \int_0^{\infty} (c_1f_1(x) + c_2f_2(x)) e^{-sx} dx \\ &= c_1 \int_0^{\infty} f_1(x)e^{-sx} dx + c_2 \int_0^{\infty} f_2(x)e^{-sx} dx \\ &= c_1\mathcal{L}[f_1](s) + c_2\mathcal{L}[f_2](s) \end{aligned}$$

Proof of (ii).

Integrating by parts one finds

$$(28.14) \quad \begin{aligned} \mathcal{L}[f'](s) &= \int_0^{\infty} e^{-st} f'(t) dt \\ &= e^{-st} f(t) \Big|_0^{\infty} - \int_0^{\infty} (-se^{-st}) f(t) dt \\ &= 0 - f(0) + s \int_0^{\infty} e^{-st} f(t) dt \\ &= s\mathcal{L}[f] - f(0) \quad . \end{aligned}$$

Similarly, (iii) is proved by integrating by parts repeatedly.