

## Homogeneous Equations with Constant Coefficients, Cont'd

Recall that the general solution of a  $2^{nd}$  order linear homogeneous differential equation

$$(17.1) \quad L[y] = y'' + p(x)y' + q(x)y = 0$$

is always a linear combination

$$(17.2) \quad y(x) = c_1y_1(x) + c_2y_2(x)$$

of two linearly independent solutions  $y_1$  and  $y_2$ , and we've seen that if we're given one solution  $y_1(x)$  we can compute a second linearly independent solution using the method of reduction of order. We will now turn to the problem of actually finding a single solution  $y_1(x)$  of (17.1).

We let us now return to the special case of a homogeneous second order linear differential equation with constant coefficients; i.e., differential equations of the form

$$(17.3) \quad y'' + py' + qy = 0$$

where  $p$  and  $q$  are constant.

We saw in Lecture 11, that one can construct solutions of the differential equation (17.3) by looking for solutions of the form

$$(17.4) \quad y(x) = e^{\lambda x} \quad .$$

Let us recall that construction. Plugging (17.4) into (17.3) yields

$$(17.5) \quad 0 = \lambda^2 e^{\lambda x} + p\lambda e^{\lambda x} + qe^{\lambda x} = (\lambda^2 + p\lambda + q) e^{\lambda x} \quad .$$

Since the exponential function  $e^{\lambda x}$  never vanishes we must have

$$(17.6) \quad \lambda^2 + p\lambda + q = 0 \quad .$$

Equation (17.6) is called the **characteristic equation** for (17.3) since for any  $\lambda$  satisfying (17.6) we will have a solution  $y(x) = e^{\lambda x}$  of (17.3).

Now because (17.6) is a quadratic equation we can employ the Quadratic Formula to find all of its roots:

$$(17.7) \quad \lambda^2 + p\lambda + q = 0 \quad \Rightarrow \quad \lambda = \frac{-p \pm \sqrt{p^2 - 4q}}{2} \quad .$$

Note that a root  $\lambda$  of (17.6) need not be a real number. Indeed, if  $p^2 - 4q < 0$ , then in order to compute  $\lambda$  via the Quadratic Formula we have to take the square root of a negative number and that forces us into the realm of complex numbers. We shall postpone temporarily the case when a root  $\lambda$  is complex and first discuss the case when the roots of (17.6) are all real. This requires  $p^2 - 4q \geq 0$ .

*Case (i):  $p^2 - 4q > 0$*

Because  $p^2 - 4q$  is positive,  $\sqrt{p^2 - 4q}$  is a positive real number and

$$(17.8) \quad \begin{aligned} \lambda_+ &= \frac{-p + \sqrt{p^2 - 4q}}{2} \\ \lambda_- &= \frac{-p - \sqrt{p^2 - 4q}}{2} \end{aligned}$$

are distinct real roots of (17.6). Thus,

$$(17.9) \quad \begin{aligned} y_1 &= e^{\lambda_+ x} \\ y_2 &= e^{\lambda_- x} \end{aligned}$$

will both be solutions of (17.3). Noting that

$$(17.10) \quad \begin{aligned} W(y_1, y_2) &= y_1 y_2' - y_1' y_2 \\ &= \lambda_- e^{\lambda_+ x} e^{\lambda_- x} - \lambda_+ e^{\lambda_+ x} e^{\lambda_- x} \\ &= (\lambda_- - \lambda_+) e^{(\lambda_+ + \lambda_-)x} \\ &= \frac{\sqrt{p^2 - 4q}}{a} e^{-\frac{p}{a}x} \end{aligned}$$

is non-zero, we conclude that if  $p^2 - 4q \neq 0$ , then the roots (17.8) furnish two linearly independent solutions of (17.3) and so the general solution is given by

$$(17.11) \quad y(x) = c_1 e^{\lambda_+ x} + c_2 e^{\lambda_- x} \quad .$$

*Case (ii):  $p^2 - 4q = 0$*

If  $p^2 - 4q = 0$ , however, this construction only gives us one distinct solution; because in this case  $\lambda_+ = \lambda_-$ . To find a second fundamental solution we must use the method of Reduction of Order.

So suppose  $y_1(x) = e^{-\frac{p}{2}x}$  is the solution corresponding to the root

$$\lambda = \frac{-p \pm \sqrt{p^2 - 4q}}{2} = \frac{-p \pm 0}{2} = \frac{-p}{2}$$

of

$$\lambda^2 + p\lambda - q = 0 \quad , \quad p^2 - 4q = 0.$$

Then the Reduction of Order formula gives us a second linearly independent solution

$$y_2(x) = y_1(x) \int^x \frac{1}{(y_1(s))^2} \exp \left[ \int^s -p(t) dt \right] ds$$

gives us a second linearly independent solution. Plugging in  $y_1(x) = e^{-\frac{p}{2}x}$  and  $p(t) = p$ , yields

$$\begin{aligned} y_2(x) &= e^{-\frac{p}{2}x} \int^x \frac{1}{(e^{-\frac{p}{2}s})^2} \exp \left[ \int^s -p dt \right] ds \\ &= e^{-\frac{p}{2}x} \int^x \frac{1}{e^{-ps}} \exp[-ps] ds \\ &= e^{-\frac{p}{2}x} \int^x e^{ps} e^{-ps} ds \\ &= e^{-\frac{p}{2}x} \int^x ds \\ &= x e^{-\frac{p}{2}x} \\ &= x y_1(x) \end{aligned}$$

In summary, for the case when  $p^2 - 4q = 0$ , we only have one root of the characteristic equation, and so we get only one distinct solution  $y_1(x)$  of the original differential equation by solving the characteristic equation for  $\lambda$ . To get a second linearly solution we must use the Reduction of Order formula; however, the result will always be the same: **the second linearly independent solution will always be  $x$  times the solution**  $y_1(x) = e^{-\frac{p}{2}x}$ . Thus, the general solution in this case will be

$$y(x) = c_1 e^{-\frac{p}{2}x} + c_2 x e^{-\frac{p}{2}x} \quad , \quad \text{if } p^2 - 4q = 0.$$

We now turn to the third and last possibility.

Case (iii):  $p^2 - 4q < 0$

In this case

$$(17.12) \quad \sqrt{p^2 - 4q}$$

will be undefined unless we introduce complex numbers. But when we set

$$(17.13) \quad \sqrt{-1} = i$$

we have

$$(17.14) \quad \sqrt{p^2 - 4q} = \sqrt{(-1)(4q - p^2)} = \sqrt{-1}\sqrt{4q - p^2} = i\sqrt{4q - p^2} \quad .$$

The square root on the right hand side is well-defined since  $4q - p^2$  is a positive number. Thus,

$$(17.15) \quad \lambda_{\pm} = \frac{-p \pm i\sqrt{4q - p^2}}{2} = \alpha \pm i\beta$$

where

$$(17.16) \quad \alpha = -\frac{p}{2} \quad , \quad \beta = \frac{\sqrt{4q - p^2}}{2} \quad ,$$

will be a complex solution of (17.6) and

$$(17.17) \quad y(x) = c_1 e^{\alpha x + i\beta x} + c_2 e^{\alpha x - i\beta x}$$

would be a solution of (17.3) if we could make sense out of the notion of an exponential function with a complex argument.

Thus, we must address the problem of ascribing some meaning to

$$(17.18) \quad e^{\alpha x + i\beta x}$$

as a function of  $x$ . To ascribe some sense to this expression we considered the Taylor series expansion of  $e^x$

$$(17.19) \quad \begin{aligned} e^x &= 1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \dots \\ &= \sum_{i=0}^{\infty} \frac{1}{i!}x^i \end{aligned}$$

Now although we do not yet understand what  $e^{\alpha x + i\beta x}$  means, we can nevertheless substitute  $\alpha x + i\beta$  for  $x$  on the right hand side of (17.19), and get a well defined series with values in the complex numbers. One can show that this series converges for all  $\alpha$ ,  $\beta$  and  $x$ . We thus take

$$(17.20) \quad e^{\alpha x + i\beta x} = \lim_{n \rightarrow \infty} \sum_{i=0}^n \frac{1}{i!} (\alpha x + i\beta x)^i$$

which agrees with (17.19) when  $\beta = 0$ .

One can also show that

$$(17.21) \quad e^{\alpha x + i\beta x} = e^{\alpha x} e^{i\beta x} .$$

Thus, when  $p^2 - 4q = 0$ , we have two complex valued solutions to (17.3)

$$(17.22) \quad y_1(x) = e^{\alpha x} e^{i\beta x} \quad \text{and} \quad y_2(x) = e^{\alpha x} e^{-i\beta x} \quad ,$$

where

$$(17.23) \quad \alpha = \frac{-p}{2} \quad , \quad \beta = \frac{\sqrt{4q - p^2}}{2} \quad .$$

A general solution of (17.3) would then be

$$(17.24) \quad y(x) = c_1 e^{\alpha x} e^{i\beta x} + c_2 e^{\alpha x} e^{-i\beta x} .$$

However, this is rarely the form in which one wants a solution of (17.3). One would prefer solutions that are **real-valued functions of  $x$**  rather than complex-valued functions of  $x$ . But these can be had as well, since if  $z = x + iy$  is a complex number, then

$$(17.25) \quad \begin{aligned} \operatorname{Re}(z) &= \frac{1}{2}(z + \bar{z}) = x \\ \operatorname{Im}(z) &= \frac{1}{2i}(z - \bar{z}) = y \end{aligned}$$

are both real numbers. Applying the Superposition Principle, it is easy to see that if

$$(17.26) \quad y(x) = e^{\alpha x} e^{i\beta x}$$

and

$$(17.27) \quad \bar{y}(x) = e^{\alpha x} e^{-i\beta x}$$

are two complex-valued solutions of (17.3), then

$$(17.28) \quad y_r(x) = \frac{1}{2}(y(x) + \bar{y}(x)) = e^{\alpha x} \left( \frac{e^{i\beta x} + e^{-i\beta x}}{2} \right)$$

and

$$(17.29) \quad y_i(x) = \frac{1}{2i}(y(x) - \bar{y}(x)) = e^{\alpha x} \left( \frac{e^{i\beta x} - e^{-i\beta x}}{2i} \right)$$

are both real-valued solutions of (17.3).

Let us now compute the series expansion of

$$(17.30) \quad \frac{e^{ix} + e^{-ix}}{2}$$

and

$$(17.31) \quad \frac{e^{ix} - e^{-ix}}{2i} .$$

$$(17.32) \quad \begin{aligned} \frac{1}{2}(e^{ix} + e^{-ix}) &= \frac{1}{2}(1 + (ix) + \frac{1}{2!}(ix)^2 + \frac{1}{3!}(ix)^3 + \dots) \\ &+ \frac{1}{2}(1 + (-ix) + \frac{1}{2!}(-ix)^2 + \frac{1}{3!}(-ix)^3 + \dots) \\ &= (1 - \frac{1}{2!}x^2 + \frac{1}{4!}x^4 + \dots) \end{aligned}$$

The expression on the right hand side is readily identified as the Taylor series expansion of  $\cos(x)$ . We thus conclude

$$(17.33) \quad \boxed{\cos(x) = \frac{e^{ix} + e^{-ix}}{2} .}$$

Similarly, one can show that

$$(17.34) \quad \boxed{\sin(x) = \frac{e^{ix} - e^{-ix}}{2i} .}$$

On the other hand, if one adds (17.33) to  $i$  times (17.34) one gets

$$(17.35) \quad \cos(x) + i \sin(x) = \frac{e^{ix} + e^{-ix}}{2} + i \frac{e^{ix} - e^{-ix}}{2i} = \frac{e^{ix} + e^{-ix} + e^{ix} - e^{-ix}}{2} = e^{ix}$$

or

$$(17.36) \quad \boxed{e^{ix} = \cos(x) + i \sin(x)}$$

Thus, the real part of  $e^{ix}$  is  $\cos(x)$ , while the pure imaginary part of  $e^{ix}$  is  $\sin(x)$ .

We now have a means of interpreting the function

$$(17.37) \quad e^{\alpha x + i\beta x}$$

in terms of elementary functions (rather than as a power series); namely,

$$(17.38) \quad e^{\alpha x + i\beta x} = e^{\alpha x} e^{i\beta x} = e^{\alpha x} (\cos(\beta x) + i \sin(\beta x)).$$

Thus,

$$(17.39) \quad \boxed{\begin{array}{l} \operatorname{Re} [e^{\alpha x + i\beta x}] = e^{\alpha x} \cos(\beta x) \quad , \\ \operatorname{Im} [e^{\alpha x + i\beta x}] = e^{\alpha x} \sin(\beta x) \quad . \end{array}}$$

I now want to show how (17.33) and (17.34) allow us to write down the general solution of a differential equation of the form

$$(17.40) \quad y'' + py' + qy = 0 \quad , \quad p^2 - 4q < 0$$

as a linear combination of real-valued functions.

Now when  $p^2 - 4q < 0$ , then

$$(17.41) \quad \lambda_{\pm} = \frac{-p \pm i\sqrt{4q - p^2}}{2} = \alpha \pm i\beta$$

are the (complex) roots of the characteristic equation

$$(17.42) \quad \lambda^2 + p\lambda + q = 0$$

corresponding to (17.40) and

$$(17.43) \quad y_{\pm}(x) = e^{\alpha x \pm i\beta x}$$

are two (complex-valued) solutions of (17.40). But since (17.40) is linear, since  $y_+$  and  $y_-$  are solutions so are

$$(17.44) \quad \begin{aligned} y_1(x) &= \frac{1}{2} (y_+(x) + y_-(x)) \\ &= \frac{1}{2} (e^{\alpha x + i\beta x} + e^{\alpha x - i\beta x}) \\ &= e^{\alpha x} \left( \frac{e^{i\beta x} + e^{-i\beta x}}{2} \right) \\ &= e^{\alpha x} \cos(\beta x) \end{aligned}$$

and

$$(17.45) \quad \begin{aligned} y_2(x) &= \frac{1}{2i} (y_+(x) - y_-(x)) \\ &= \frac{1}{2i} (e^{\alpha x + i\beta x} - e^{\alpha x - i\beta x}) \\ &= e^{\alpha x} \left( \frac{e^{i\beta x} - e^{-i\beta x}}{2i} \right) \\ &= e^{\alpha x} \sin(\beta x) \quad . \end{aligned}$$

Note that  $y_1$  and  $y_2$  are both **real-valued functions**.

We conclude that if the characteristic equation corresponding to

$$(17.46) \quad y'' + py' + qy = 0$$

has two complex roots

$$(17.47) \quad \lambda = \alpha \pm i\beta$$

then the general solution is

$$(17.48) \quad \boxed{y(x) = c_1 e^{\alpha x} \cos(\beta x) + c_2 e^{\alpha x} \sin(\beta x) \quad .}$$

EXAMPLE 17.1. The differential equation

$$(17.49) \quad y'' - 2y' - 3y$$

has as its characteristic equation

$$(17.50) \quad \lambda^2 - 2\lambda - 3 = 0 \quad .$$

The roots of the characteristic equation are given by

$$(17.51) \quad \begin{aligned} \lambda &= \frac{2 \pm \sqrt{4+12}}{2} \\ &= 3, -1 \quad . \end{aligned}$$

These are distinct real roots, so the general solution is

$$(17.52) \quad y(x) = c_1 e^{3x} + c_2 e^{-x} \quad .$$

EXAMPLE 17.2. The differential equation

$$(17.53) \quad y'' + 4y' + 4y = 0$$

has

$$(17.54) \quad \lambda^2 + 4\lambda + 4 = 0$$

as its characteristic equation. The roots of the characteristic equation are given by

$$(17.55) \quad \begin{aligned} \lambda &= \frac{-4 \pm \sqrt{16-16}}{2} \\ &= -2 \quad . \end{aligned}$$

Thus we have a double root and the general solution is

$$(17.56) \quad y(x) = c_1 e^{-2x} + c_2 x e^{-2x} \quad .$$

EXAMPLE 17.3. The differential equation

$$(17.57) \quad y'' + y' + y = 0$$

has

$$(17.58) \quad \lambda^2 + \lambda + 1 = 0$$

as its characteristic equation. The roots of the characteristic equation are

$$(17.59) \quad \begin{aligned} \lambda &= \frac{-1 \pm \sqrt{1-4}}{2} \\ &= -\frac{1}{2} \pm i \frac{\sqrt{3}}{2} \end{aligned}$$

and so the general solution is

$$(17.60) \quad y(x) = c_1 e^{-\frac{1}{2}x} \cos\left(\frac{\sqrt{3}}{2}x\right) + c_2 e^{-\frac{1}{2}x} \sin\left(\frac{\sqrt{3}}{2}x\right) \quad .$$