

Convex-Ear Decompositions and the Flag h-Vector

Jay Schweig

University of Kansas
Kansas, U.S.A.

jschweig@math.ku.edu

Submitted: Jun 10, 2010; Accepted: Dec 13, 2010; Published: Jan 5, 2011

Mathematics Subject Classification: 05E45, 06F30, 52B22

Abstract

We prove a theorem allowing us to find convex-ear decompositions for rank-selected subposets of posets that are unions of Boolean sublattices in a coherent fashion. We then apply this theorem to geometric lattices and face posets of shellable complexes, obtaining new inequalities for their h-vectors. Finally, we use the latter decomposition to give a new interpretation to inequalities satisfied by the flag h-vectors of face posets of Cohen-Macaulay complexes.

1 Introduction

The *f*-vector of a finite simplicial complex Δ , which counts the number of faces of the complex in each dimension, is arguably its most fundamental invariant. The *h*-vector of Δ is the image of its f-vector under an invertible transformation. Somewhat surprisingly, properties of a complex's f-vector are sometimes better expressed through its h-vector. A good example of this phenomenon are the Dehn-Sommerville relations (see, for instance, [17]), which state that the h-vector of a simplicial polytope boundary is symmetric.

The main complexes we study in this paper are all *order complexes*, namely complexes whose simplices correspond to chains in posets. Since a poset and its order complex hold the same information, we often refer to them interchangeably. E.g., we may speak of the *facets* or *h*-vector of a poset, or to a *chain* in an order complex.

Convex-ear decompositions were first introduced by Chari in [3]. Heuristically, a complex admits a convex-ear decomposition if it is a union of simplicial polytope boundaries which fit together coherently (see Definition 2.3). Suppose a $(d - 1)$ -dimensional complex Δ admits such a decomposition. In [3], Chari shows that the h-vector (h_0, h_1, \dots, h_d) of Δ satisfies, for $i < d/2$, $h_i \leq h_{i+1}$ and $h_i \leq h_{d-i}$. In [15], Swartz shows that Δ is 2-CM, and that $(h_0, h_1 - h_0, h_2 - h_1, \dots, h_{\lfloor d/2 \rfloor} - h_{\lfloor d/2 \rfloor - 1})$ is an M-vector (called an *O*-sequence by some authors). Convex-ear decompositions have proven quite useful, as they have been applied to coloop-free matroid complexes [3], geometric lattices [9], coloring complexes [6],

d -divisible partition lattices [16], coset lattices of relatively complemented finite groups [16], and finite buildings [15].

In [11], we find a convex-ear decomposition for rank-selected subposets of supersolvable lattices with nowhere-zero Möbius functions. In the process, we obtain a decomposition for order complexes of rank-selected subposets of the Boolean lattice B_d . In this paper we build upon this result with Theorem 3.1, which gives convex-ear decompositions for rank-selected subposets of posets that are unions of Boolean lattices, pieced together nicely. In Section 3 we recall several useful results from [11], and then prove this theorem.

Sections 4 and 5 are devoted to the applications of Theorem 3.1 to geometric lattices and face posets of shellable complexes, respectively. Taken together, these results (along with those of Chari and Swartz) give us the following.

Theorem 1.1. *Let Δ be the order complex of a rank-selected subposet of either*

1: a geometric lattice, or

2: the face poset of the codimension-1 skeleton of a Cohen-Macaulay complex.

Then Δ is 2-CM and its h -vector (h_0, h_1, \dots, h_d) satisfies, for all $i < d/2$, $h_i \leq h_{i+1}$ and $h_i \leq h_{d-i}$. Moreover, $(h_0, h_1 - h_0, h_2 - h_1, \dots, h_{\lfloor d/2 \rfloor} - h_{\lfloor d/2 \rfloor - 1})$ is an M -vector.

For item 2 above, we use the fact that the set of h -vectors in question remains unchanged if we replace the Cohen-Macaulay requirement with the stronger condition of shellability (see the proof of Theorem 6.2).

Finally, in Section 6, we use the decomposition from Section 5 and techniques similar to those in [9] to prove that the flag h -vector $\{h_S\}$ of a Cohen-Macaulay complex's face poset satisfies $h_T \leq h_S$ whenever S dominates T (in the sense of Definition 2.10).

2 Preliminaries

We assume a familiarity with simplicial complexes and partially ordered sets (see [14]). All our simplicial complexes will be finite and pure.

The f -vector of a $(d - 1)$ -dimensional simplicial complex Δ is the integral sequence (f_0, f_1, \dots, f_d) , where f_i counts the number of $(i - 1)$ -dimensional faces of Δ . The h -vector of Δ is the sequence (h_0, h_1, \dots, h_d) given by

$$\sum_{i=0}^d f_i(t - 1)^{d-i} = \sum_{i=0}^d h_i t^{d-i}$$

We use the following alternate definition of shellability, easily seen to be equivalent to the standard one (see [1]).

Definition 2.1. *Let F_1, F_2, \dots, F_t be an ordering of the facets of Δ . This ordering is a shelling if and only if for all $j < k$ there exists a $j' < k$ satisfying*

$$F_j \cap F_k \subseteq F_{j'} \cap F_k = F_{j'} - x$$

for some element x of $F_{j'}$.

We also use the following result of Danaraj and Klee for showing that a given complex is a ball.

Theorem 2.2 ([4]). *Let Δ be a full-dimensional shellable proper subcomplex of a sphere. Then Δ is a ball.*

2.1 Convex-ear decompositions

Definition 2.3. *Let Δ be a $(d-1)$ -dimensional complex. We say Δ admits a convex-ear decomposition, or c.e.d., if there exists a sequence of pure $(d-1)$ -dimensional subcomplexes $\Sigma_1, \Sigma_2, \dots, \Sigma_t$ such that*

$$i: \bigcup_{i=1}^t \Sigma_i = \Delta.$$

ii: *For $i > 1$, Σ_i is a proper subcomplex of the boundary complex of a d -dimensional simplicial polytope, while Σ_1 is the boundary complex of a d -dimensional simplicial polytope.*

iii: *Each Σ_i , for $i > 1$ is a topological ball.*

$$iv: \text{For } i > 1, \Sigma_i \cap \left(\bigcup_{j=1}^{i-1} \Sigma_j\right) = \partial\Sigma_i.$$

Convex-ear decompositions were introduced by Chari in [3], where the following was proven.

Theorem 2.4. *When Δ admits a convex-ear decomposition its h -vector satisfies, for all $i \leq \lfloor d/2 \rfloor$,*

$$1: h_i \leq h_{i+1} \text{ and}$$

$$2: h_i \leq h_{d-i}.$$

In [15], Swartz proved the following analogue of the g -Theorem for complexes with convex-ear decompositions. An M -vector is the degree sequence of an order ideal of monomials.

Theorem 2.5. *Let Δ be a complex admitting a convex-ear decomposition, with h -vector (h_0, h_1, \dots, h_d) . Then the vector*

$$(h_0, h_1 - h_0, h_2 - h_1, \dots, h_{\lfloor d/2 \rfloor} - h_{\lfloor d/2 \rfloor - 1})$$

is an M -vector. Furthermore, Δ is 2-CM.

2.2 Order complexes and flag vectors

Recall that the *order complex* of a poset P , which we write as $\Delta(P)$, is the simplicial complex whose faces are chains in P . If P has a unique minimal element $\hat{0}$ or a unique maximal element $\hat{1}$, we do not include these in the order complex. That is, simplices in $\Delta(P)$ are chains in $P - \{\hat{0}, \hat{1}\}$. All our posets are ranked.

For the remainder of this section, let P be a rank- d poset with a $\hat{0}$ and $\hat{1}$. A *labeling* of P is a function $\lambda : \{(x, y) \in P^2 : y \text{ covers } x\} \rightarrow \mathbb{Z}$. For a saturated chain

$$\mathbf{c} := x = x_0 < x_1 < \cdots < x_t = y$$

the λ -label of \mathbf{c} , written $\lambda(\mathbf{c})$, is the word

$$\lambda(x_0, x_1)\lambda(x_1, x_2) \cdots \lambda(x_{t-1}, x_t).$$

Definition 2.6. A labeling λ of P is an EL-labeling if:

- 1: in each interval $[x, y]$ in P , there is a unique saturated chain with a strictly increasing label, and
- 2: the label of this chain is lexicographically first among the labels of all saturated chains in $[x, y]$.

If λ is an EL-labeling of P and each maximal chain is labeled with a permutation of $[d]$ (that is, an element of \mathcal{S}_d) then λ is called an \mathcal{S}_d -EL-labeling.

Example 2.7. The Boolean lattice B_d admits an \mathcal{S}_d -EL-labeling in an obvious way: if y covers x then $y = x \cup \{i\}$, so set $\lambda(x, y) = i$.

When P admits an EL-labeling λ and \mathbf{c} is a chain in P , we write $\Upsilon_\lambda(\mathbf{c})$ to denote the maximal chain of P obtained by filling in each gap in \mathbf{c} with the unique chain in that interval with increasing λ -label. EL-labelings were introduced by Björner and Wachs, where the following was shown.

Theorem 2.8 ([2]). If P admits an EL-labeling then $\Delta(P)$ is shellable.

For any $S \subseteq [d-1]$ and any maximal chain

$$\mathbf{c} := \hat{0} = x_0 < x_1 < x_2 < \cdots < x_d = \hat{1}$$

of P , let \mathbf{c}_S denote the chain of elements of \mathbf{c} whose ranks lie in $S \cup \{0, d\}$. The *rank-selected subposet* P_S is the subposet of P whose maximal chains are all of the form \mathbf{c}_S , as \mathbf{c} ranges over all maximal chains of P . Equivalently, P_S is the poset P restricted to all elements with ranks in $S \cup \{0, d\}$.

For any $S \subseteq [d-1]$, let f_S be the number of maximal chains in P_S . The collection $\{f_S\}$ is known as the *flag f-vector* of P . Note that the flag f-vector of P refines its f-vector, as clearly

$$f_i(P) = \sum_{S \subseteq [d-1], |S|=i} f_S(P).$$

The *flag h-vector* of P is the collection $\{h_S\}$ defined by

$$h_S = \sum_{T \subseteq S} (-1)^{|S-T|} f_T.$$

By inclusion-exclusion, the above is equivalent to $f_T = \sum_{S \subseteq T} h_S$. It follows that the h-vector is refined by the flag h-vector, namely

$$h_i(P) = \sum_{S \subseteq [d-1], |S|=i} h_S(P).$$

When P has an EL-labeling, its flag h-vector has a nice enumerative interpretation.

Theorem 2.9 ([2]). *The flag h-vector $\{h_S\}$ of a poset P with EL-labeling λ is given as follows: h_S counts the number of maximal chains of P whose λ -labels have descent set S .*

2.3 Dominance in \mathcal{S}_d .

Let σ be a permutation in the symmetric group \mathcal{S}_d . We view σ as a word in $[d]$, writing $\sigma = \sigma(1)\sigma(2) \cdots \sigma(d)$. If $\sigma(i) < \sigma(i+1)$ call the interchanging of $\sigma(i)$ and $\sigma(i+1)$ in σ a *switch*.

Recall that the *weak order* on \mathcal{S}_d , for which we write $<_w$, is the partial order given by the following property: $\sigma <_w \tau$ if and only if τ can be obtained from σ by a sequence of switches.

For $S \subseteq [d-1]$, let D_S^d denote the set of permutation in \mathcal{S}_d whose descent sets equal S :

$$\sigma \in D_S^d \Leftrightarrow \{i : \sigma(i) > \sigma(i+1)\} = S.$$

Definition 2.10. *Let $S, T \subseteq [d-1]$. We say that S dominates T if there exists an injection $\phi : D_T^d \rightarrow D_S^d$ such that $\sigma <_w \phi(\sigma)$ for all $\sigma \in D_T^d$.*

For example, let $d = 4$. Then the set $\{1, 3\}$ dominates the set $\{1\}$ via the map

$$\sigma(1)\sigma(2)\sigma(3)\sigma(4) \rightarrow \sigma(1)\sigma(2)\sigma(4)\sigma(3).$$

For a further discussion of dominance in the symmetric group, see [9] or [5].

3 A decomposition theorem

The goal of this section is to prove the following theorem, which we will apply in Sections 4 and 5.

Theorem 3.1. *Let P be a rank- d poset with a $\hat{0}$ and a $\hat{1}$, and suppose P_1, P_2, \dots, P_r are subposets of P satisfying the following properties.*

- 1: *Each P_i is isomorphic to the Boolean lattice B_d .*

2: Every chain in P is a chain in some P_i . Equivalently,

$$\Delta(P) = \bigcup_{i=1}^r \Delta(P_i).$$

3: Each P_i has an \mathcal{S}_d -EL-labeling λ_i with the following property: if \mathbf{e} is a chain in P_i that is also a chain in some P_j for $j < i$, then $\Upsilon_{\lambda_i}(\mathbf{e})$ is a chain in P_j for some $j < i$.

Then $\Delta(P_S)$ admits a convex-ear decomposition for any $S \subseteq [d-1]$.

The proof of this theorem relies heavily on our work from [11], where a c.e.d. of $\Delta((B_d)_S)$ was given for any $S \subseteq [d-1]$. We now review some results from [11] which will be helpful in proving Theorem 3.1.

Fix some $S \subseteq [d-1]$ and an \mathcal{S}_d -EL-labeling λ of B_d . Let $\mathbf{d}_1, \mathbf{d}_2, \dots, \mathbf{d}_t$ be all maximal chains in B_d whose λ -labels have descent set S , written in lexicographic order of their λ -labels. For each i , let L_i be the subposet of $(B_d)_S$ generated by the set of maximal chains

$$\{\mathbf{c}_S : \mathbf{c} \text{ is a maximal chain of } B_d \text{ with } \mathbf{c}_{[d-1] \setminus S} = (\mathbf{d}_i)_{[d-1] \setminus S}\}$$

Finally, let Γ_i be the simplicial complex with facets given by maximal chains in L_i that are not chains in L_j for any $j < i$.

Theorem 3.2 ([11]). *The sequence of complexes $\Gamma_1, \Gamma_2, \dots, \Gamma_t$ is a convex-ear decomposition of $\Delta((B_d)_S)$.*

The following lemmata, whose proofs we omit, are shown in [11].

Lemma 3.3. *Let \mathbf{e} be a maximal chain of some L_i . Then \mathbf{e} is a facet of Γ_i if and only if*

$$(\Upsilon_{\lambda}(\mathbf{e}))_{[d-1] \setminus S} = (\mathbf{d}_i)_{[d-1] \setminus S}.$$

Lemma 3.4. *Let \mathbf{c} be the unique maximal chain of B_d with increasing λ -label. Then*

$$\Upsilon_{\lambda}((\mathbf{d}_1)_{[d-1] \setminus S}) = \mathbf{c}.$$

Lemma 3.5. *Let $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_m$ be all maximal chains corresponding to facets of Γ_i , ordered so that $\lambda(\Upsilon_{\lambda}(\mathbf{e}_k))$ lexicographically precedes $\lambda(\Upsilon_{\lambda}(\mathbf{e}_j))$ whenever $j < k$. Then for all j and k with $j < k$, there exists a $j' < k$ satisfying*

$$\mathbf{e}_j \cap \mathbf{e}_k \subseteq \mathbf{e}_{j'} \cap \mathbf{e}_k = \mathbf{e}_{j'} - x$$

for some element x of $\mathbf{e}_{j'}$.

Lemma 3.5, together with Theorem 2.2 and Definition 2.1, proves that Γ_i is a topological ball for $i > 1$. We are now ready to prove our main theorem.

Proof of Theorem 3.1. The basic idea is to iterate the decomposition provided by Theorem 3.2. Indeed, Theorem 3.2 gives us a c.e.d. of $\Delta((P_1)_S)$. Now suppose we have a c.e.d. for $X = \bigcup_{i=1}^{q-1} \Delta((P_i)_S)$ for some q with $2 \leq q \leq r$. We show that we can extend this to a c.e.d. of $X \cup \Delta((P_q)_S)$. For ease of notation, let $\lambda = \lambda_q$ and $\Upsilon = \Upsilon_\lambda$.

Taking our cue from the decomposition of $(B_d)_S$ described above, let $\mathbf{d}_1, \mathbf{d}_2, \dots, \mathbf{d}_t$ be all maximal chains of P_q whose λ -labels have descent set S , and let the above order be such that $\lambda(\mathbf{d}_i)$ lexicographically precedes $\lambda(\mathbf{d}_j)$ for $i < j$. For each i , define L_i and Γ_i as in Theorem 3.2. Finally, let Σ_i be the simplicial complex whose facets are maximal chains in Γ_i that are not chains in X . We claim that the sequence $\Sigma_1, \Sigma_2, \dots, \Sigma_t$ (once we remove all $\Sigma_i = \emptyset$) extends the c.e.d. of X . We prove each property of Definition 2.3 separately.

By definition, each $\Gamma_i \subseteq \Sigma_i \cup X$. Since $\bigcup_{i=1}^t \Gamma_i = \Delta((P_q)_S)$ (by Theorem 3.2), $X \cup (\bigcup_{i=1}^t \Sigma_i) = X \cup \Delta((P_q)_S)$, so property (i) holds.

Property (ii) is easily verified as well. Since each Γ_i for $i > 1$ is a proper subcomplex of a simplicial polytope boundary, so is each $\Sigma_i \subseteq \Gamma_i$. However, as Γ_1 is a simplicial polytope boundary, we need to show that the inclusion $\Sigma_1 \subseteq \Gamma_1$ is proper. This follows from Lemma 3.4, which says that \mathbf{c}_S is a facet of Γ_1 , where \mathbf{c} is the unique maximal chain in P_q with increasing λ -label. Because $\mathbf{c} = \Upsilon(\hat{0} < \hat{1})$ and $\hat{0} < \hat{1}$ is a chain in all P_i , it follows that \mathbf{c}_S is not a facet of Σ_1 .

Now fix $i \geq 1$. To prove property (iii), we employ the techniques (and notation) of Lemma 3.5. Let $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_m$ be all maximal chains of Σ_i , ordered so that $\lambda(\Upsilon(\mathbf{e}_k))$ precedes $\lambda(\Upsilon(\mathbf{e}_j))$ whenever $j < k$. Now choose some j and k with $j < k$. Because $\Sigma_i \subseteq \Gamma_i$, Lemma 3.5 produces a maximal chain \mathbf{e} in Γ_i satisfying $\mathbf{e}_j \cap \mathbf{e}_k \subseteq \mathbf{e} \cap \mathbf{e}_k = \mathbf{e} - x$ for some element x of \mathbf{e} , with $\lambda(\Upsilon(\mathbf{e}_k))$ lexicographically preceding $\lambda(\Upsilon(\mathbf{e}))$. To finish the proof, we just need to show that \mathbf{e} is a facet of Σ_i . That is, we need to show that \mathbf{e} is not a chain in X .

By Lemma 3.3, $\mathbf{e} \cap \mathbf{e}_k = \mathbf{e} - x$ implies that $\Upsilon(\mathbf{e}) \cap \Upsilon(\mathbf{e}_k) = \Upsilon(\mathbf{e}) - x$. Because P_q is a Boolean lattice, it has exactly two maximal chains containing $\Upsilon(\mathbf{e}) - x$ as a subchain. Hence, these chains must be $\Upsilon(\mathbf{e})$ and $\Upsilon(\mathbf{e}_k)$. Because $\lambda(\Upsilon(\mathbf{e}_k))$ precedes $\lambda(\Upsilon(\mathbf{e}))$, we have $\Upsilon(\mathbf{e}_k) = \Upsilon(\Upsilon(\mathbf{e}) - x)$. If \mathbf{e} were a chain in X , then $\Upsilon(\mathbf{e}) - x$ would be as well. But then, since \mathbf{e}_k is a subchain of $\Upsilon(\Upsilon(\mathbf{e}) - x)$, we would have that \mathbf{e}_k is in X , a contradiction.

For property (iv) consider some i , and note that a chain \mathbf{e} is in $\partial\Sigma_i$ if and only if there exist two maximal chains \mathbf{e}_{old} and \mathbf{e}_{new} , each containing \mathbf{e} as a subchain, such that \mathbf{e}_{old} is a chain in $X \cup (\bigcup_{j=1}^{i-1} \Sigma_j)$ and \mathbf{e}_{new} is a chain in Σ_i . Thus,

$$\partial\Sigma_i \subseteq \Sigma_i \cap \left(X \cup \left(\bigcup_{j=1}^{i-1} \Sigma_j \right) \right).$$

To prove the reverse inclusion, let \mathbf{e} be a non-maximal chain in $\Sigma_i \cap (X \cup (\bigcup_{j=1}^{i-1} \Sigma_j))$. Then by definition \mathbf{e} must be a subchain of some maximal chain in Σ_i , and we can take this chain to be \mathbf{e}_{new} . To find \mathbf{e}_{old} , we consider two cases. First, if \mathbf{e} is a chain of Σ_j for some $j < i$, then by Theorem 3.2 there must be some maximal chain \mathbf{e}_{old} of Σ_j for some $j < i$. Second, if \mathbf{e} is a chain in X , then $\Upsilon(\mathbf{e})$ must be in X as well. Setting $\mathbf{e}_{\text{old}} = (\Upsilon(\mathbf{e}))_S$

completes the proof of property (iv).

Thus, we can extend the c.e.d. of X to one of $X \cup \Delta((P_q)_S)$. Continuing in this fashion, we get a c.e.d. of

$$\bigcup_{i=1}^r \Delta((P_i)_S).$$

By hypothesis every chain in P is a chain in some P_i and the above union equals $\Delta(P_S)$, proving the theorem. \square

4 Rank-selected geometric lattices

We first apply Theorem 3.1 to geometric lattices. We assume a basic familiarity with matroid theory, including the cryptomorphism between matroids and geometric lattices. For background, see [1] or [10].

Let P be a rank- d geometric lattice. In [9], Nyman and Swartz show that $\Delta(P)$ admits a convex-ear decomposition. We open this section by briefly describing their technique.

Let a_1, a_2, \dots, a_ℓ be a fixed linear ordering of the atoms of P . The *minimal labeling* ν of P is defined as follows: if y covers x , then $\nu(x, y) = \min\{i : x \vee a_i = y\}$. We view P as the lattice of flats of a simple matroid M .

Lemma 4.1 ([1]). *The minimal labeling ν is an EL-labeling.*

Lemma 4.2 ([1]). *Suppose the ν -label of a maximal chain \mathbf{c} of P is a word in some subset $B \subseteq \{a_1, a_2, \dots, a_\ell\}$. Then B is an nbc-basis of M .*

Now let B_1, B_2, \dots, B_t be all the nbc-bases of M listed in lexicographic order. Fix some $j \leq t$ and let $B_j = \{a_{i_1}, a_{i_2}, \dots, a_{i_d}\}$ with $i_1 < i_2 < \dots < i_d$. For a permutation $\sigma \in \mathcal{S}_d$, define a maximal chain \mathbf{c}_σ^j of P by

$$\mathbf{c}_\sigma^j := \hat{0} < a_{i_{\sigma(1)}} < (a_{i_{\sigma(1)}} \vee a_{i_{\sigma(2)}}) < \dots < (a_{i_{\sigma(1)}} \vee a_{i_{\sigma(2)}} \vee \dots \vee a_{i_{\sigma(d)}}).$$

The *basis labeling* $\lambda_j(\mathbf{c}_\sigma^j)$ of \mathbf{c}_σ^j is the word $i_{\sigma(1)}i_{\sigma(2)} \dots i_{\sigma(d)}$.

For each i with $1 \leq i \leq t$, let P_i be the subposet of P whose set of maximal chains is $\{\mathbf{c}_\sigma^i : \sigma \in \mathcal{S}_d\}$ and let Σ_i be the simplicial complex whose facets are maximal chains in P_i that are not chains in P_j for any $j < i$.

Theorem 4.3 ([9]). $\Sigma_1, \Sigma_2, \dots, \Sigma_t$ is a convex-ear decomposition of $\Delta(P)$.

The next lemma, shown in [9], is the key tool in proving the above theorem.

Lemma 4.4. *A chain \mathbf{c} in P_i is in Σ_i if and only if*

$$\lambda_i(\mathbf{c}) = \nu(\mathbf{c}).$$

The main theorem of this section is the following:

Theorem 4.5. *Let P be a rank- d geometric lattice. Then $\Delta(P_S)$ admits a convex-ear decomposition for any $S \subseteq [d - 1]$.*

Proof. We show that P satisfies the hypotheses of Theorem 3.1, proving each of the three properties separately.

First note that each P_i is isomorphic to the Boolean lattice B_d under the mapping $a_{i_{\sigma(1)}} \vee a_{i_{\sigma(2)}} \vee \dots \vee a_{i_{\sigma(m)}} \rightarrow \{\sigma(1), \sigma(2), \dots, \sigma(m)\}$, and so property (1) holds. Moreover, the basis labeling λ_i is an \mathcal{S}_d -EL-labeling of P_i (though with the alphabet $\{i_1, i_2, \dots, i_d\}$ rather than $[d]$).

By Lemma 4.2, the ν -label of any maximal chain \mathbf{c} is a word in some nbc-basis (say B_i). Thus \mathbf{c} is a chain in P_i , meaning property (2) holds.

To show property (3), fix some i and suppose that \mathbf{e} is a non-maximal chain in P_i that is also a chain in P_j for some $j < i$. Suppose that j is the least such integer, and consider the maximal chain $\mathbf{c} = \Upsilon_{\lambda_j}(\mathbf{e})$. This chain can clearly not be in L_k for any $k < j$, because then \mathbf{e} would be a chain in L_k , contradicting the minimality of j . Thus $\lambda_j(\mathbf{c}) = \nu(\mathbf{c})$ by Lemma 4.4, meaning $\mathbf{c} = \Upsilon_{\nu}(\mathbf{e})$. Now consider the chain $\mathbf{c}' = \Upsilon_{\lambda_i}(\mathbf{e})$. If \mathbf{c}' is not a chain in L_k for any $k < i$ then, again by Lemma 4.4, $\nu(\mathbf{c}') = \lambda_i(\mathbf{c}')$. But then $\mathbf{c}' = \Upsilon_{\nu}(\mathbf{e})$, which is a contradiction since the chain $\Upsilon_{\nu}(\mathbf{e})$ is uniquely determined. Thus $\Upsilon_{\lambda_i}(\mathbf{e})$ must be a chain in L_k for some $k < i$. Applying Theorem 3.1 completes the proof. \square

5 Rank-selected face posets

The main result of this section can be seen as motivated by Hibi's result ([7]) that the codimension-1 skeleton of a shellable complex Σ is 2-Cohen-Macaulay. For a simplicial complex Σ we write P_{Σ} to mean its *face poset*, the poset of all faces of Σ ordered by inclusion. Note that P_{Σ} usually does not have a unique maximal element, but the notion of the rank-selected subposet $(P_{\Sigma})_S$ easily generalizes.

Theorem 5.1. *Let Σ be a $(d - 1)$ -dimensional shellable complex. Then $\Delta((P_{\Sigma})_S)$ admits a convex-ear decomposition for any $S \subseteq [d - 1]$.*

Proof. We wish to apply Theorem 3.1 but, as noted above, a slight adjustment is needed: unless Σ consists of a single facet, P_{Σ} has no maximal element. To this end, let P be the poset P_{Σ} with all its maximal elements identified. As usual, let $\hat{1}$ denote the maximal element of P . Clearly, for any $S \subseteq [d - 1]$,

$$\Delta(P_S) = \Delta((P_{\Sigma})_S).$$

So, it suffices to apply Theorem 3.1 to P . Fix a shelling F_1, F_2, \dots, F_r of Σ , and for each i let P_i be the face poset of F_i (but with its maximal element F_i replaced with $\hat{1}$, the maximal element of P). We claim that the sequence P_1, P_2, \dots, P_r satisfies the hypotheses of Theorem 3.1. Property (1) follows immediately, as the face poset of a $(d - 1)$ -dimensional simplex is isomorphic to B_d .

For property (2), let \mathbf{c} be a maximal chain of P , and let x be its element of rank $d - 1$. Then x is a face of some facet F_i , meaning P_i contains the chain \mathbf{c} .

The proof of property (3) relies on the following fact, whose proof is immediate.

Fact 5.2. *Let \mathbf{e} be a non-maximal chain in some P_i , and let x be its element of highest rank $\neq d$. Then \mathbf{e} is not a chain in P_j for any $j < i$ if and only if, when viewed as a face of F_i , x contains the unique minimal new face $r(F_i)$.*

Now fix some i , and let V be the set of vertices of the facet F_i . Any bijection $\phi : V \rightarrow [d]$ induces an \mathcal{S}_d -EL-labeling λ_ϕ of P_i in the obvious way: For $x, y \in P_i$ with $y = x \cup \{v\}$ for some vertex v of F_i , set $\lambda_\phi(x, y) = \phi(v)$ (if $y = \hat{1}$, let $\lambda_\phi(x, y)$ be the sole vertex in $V \setminus x$). Let $\phi : F_i \rightarrow [d]$ be any bijection that labels vertices in $r(F_i)$ last. That is, if $v \in r(F_i)$ and $w \in F_i \setminus r(F_i)$ then $\phi(w) < \phi(v)$. Set $\lambda_i = \lambda_\phi$. Now suppose \mathbf{e} is a non-maximal chain in P_i that is also a chain in P_j for some $j < i$, and let x be the element of \mathbf{e} of highest rank $\neq d$. By Fact 5.2, $r(F_i) \not\subseteq x$. If v is the vertex in $F_i \setminus x$ with the greatest ϕ -label then, by definition of ϕ , $v \in r(F_i)$. Letting y be the element of $\Upsilon_{\lambda_i}(\mathbf{e})$ of rank $d - 1$, it follows that $v \notin y$. So $\Upsilon_{\lambda_i}(\mathbf{e})$ is a chain in P_k for some $k < i$, and property (3) holds. \square

In many cases, the above theorem does not hold if $d \in S$. For example, if Σ is the shellable complex consisting of two 2-dimensional simplices joined at a common boundary facet and $S = \{2, 3\} \subseteq [3]$, then $\Delta((P_\Sigma)_S)$ does not admit a c.e.d., as it is a tree. We can, however, conjecture the following.

Conjecture 5.3. *When Σ is a $(d - 1)$ -dimensional complex admitting a convex-ear decomposition and $S \subseteq [d]$, the complex $\Delta((P_\Sigma)_S)$ admits a convex-ear decomposition.*

If $d \notin S$, the above conjecture follows from Theorem 5.1, so we need only consider cases where $d \in S$.

Example 5.4. *Let Δ be a triangulation of the dunce cap, and let P_Δ be its face poset. Then $(P_\Delta)_S$ admits a convex-ear decomposition for any $S \subseteq [3]$ with $|S| = 2$. To see this, note that a simple graph (viewed as a 1-dimensional simplicial complex) has a c.e.d. if and only if it is 2-connected.*

It is well known that Δ is Cohen-Macaulay, and it is easy to see that Δ is not 2-Cohen-Macaulay. Thus, the converse to Theorem 5.1 is false.

Now recall that a $(d - 1)$ -dimensional complex Σ with vertex set V is called *balanced* if there exists a $\psi : V \rightarrow [d]$ such that $\psi(v) \neq \psi(w)$ whenever v and w are in a common face of Σ . The function ψ is called a *coloring* of Σ .

The order complex of any graded poset P is always balanced: For a vertex v of $\Delta(P)$, simply let $\psi(v)$ be the rank of v when considered as an element of the poset P . Thus the barycentric subdivision of any simplicial complex is balanced, since it is the order complex of its face poset.

If Σ is a $(d - 1)$ -dimensional balanced complex with coloring ψ and $S \subseteq [d]$, define Σ_S to be the subcomplex of Σ with faces $\{F \in \Sigma : \psi(v) \in S \text{ for all } v \in F\}$. With these new definitions, we can rephrase Theorem 5.1 in a more geometric tone.

Corollary 5.5. *Let Σ' be a $(d - 1)$ -dimensional shellable complex, and let Σ be the first barycentric subdivision of its codimension-1 skeleton. Then, for any coloring ψ of the vertices of Σ and any $S \subseteq [d - 1]$, the complex Σ_S admits a convex-ear decomposition.*

6 The flag h-vector of a face poset

Our goal in this section is to prove an analogue of the following theorem, shown in [9], for face posets of Cohen-Macaulay complexes. In this section, we assume a basic working knowledge of the Stanley-Reisner ring of a simplicial complex and its Hilbert series (see [13]).

Theorem 6.1 ([9]). *Let L be a rank- d geometric lattice, and let $S, T \subseteq [d - 1]$. If S dominates T , the flag h-vector of $\Delta(L)$ satisfies $h_T \leq h_S$.*

We state the main theorem now, but postpone its proof until after Theorem 6.5.

Theorem 6.2. *Let K be a d -dimensional Cohen-Macaulay simplicial complex with face poset P , and let $\Delta = \Delta(P)$. Let $S, T \subseteq [d - 1]$, and suppose that S dominates T . Then the flag h-vector of Δ satisfies $h_T \leq h_S$.*

Lemma 6.3. *Let P be a rank- d poset with a $\hat{0}$ and $\hat{1}$ whose order complex $\Delta = \Delta(P)$ is a ball. Let Δ' be the set of faces in $\Delta - \partial\Delta - \emptyset$, let $\{f'_S\}$ be the flag f-vector of Δ' , and let $\{h_S\}$ be the flag h-vector of Δ . Then*

$$\sum_{S \subseteq [d-1]} f'_S \prod_{i \notin S} (\nu_i - 1) = \sum_{S \subseteq [d-1]} h_{[d-1]-S} \prod_{i \notin S} \nu_i$$

Proof. Under the fine grading of the face ring $k[\Delta]$, $F(k[\Delta], \lambda) = \sum_{F \in \Delta} \prod_{x_i \in F} \frac{\lambda_i}{1 - \lambda_i}$. We specialize this grading to accommodate the flag h-vector as follows: identify λ_i and λ_j whenever the vertices in Δ to which they correspond have the same rank r (as elements of P). Call this new variable ν_r . This specialized grading yields:

$$F(k[\Delta], \nu) = \sum_{S \subseteq [d-1]} f_S \prod_{i \in S} \frac{\nu_i}{1 - \nu_i}$$

We put this over the common denominator of $\prod_{i \in [d-1]} (1 - \nu_i)$ to obtain:

$$F(k[\Delta], \nu) = \sum_{S \subseteq [d-1]} \frac{f_S \prod_{i \in S} \nu_i \prod_{i \notin S} (1 - \nu_i)}{\prod_{i \in [d-1]} (1 - \nu_i)} = \sum_{S \subseteq [d-1]} \frac{h_S \prod_{i \in S} \nu_i}{\prod_{i \in [d-1]} (1 - \nu_i)} \quad (1)$$

The following equation is Corollary II.7.2 from [13] (note that Δ is $(d - 2)$ -dimensional):

$$(-1)^{d-1} F(k[\Delta], 1/\lambda) = (-1)^{d-2} \tilde{\chi}(\Delta) + \sum_{F \in \Delta'} \prod_{x_i \in F} \frac{\lambda_i}{1 - \lambda_i}$$

Noting that $\tilde{\chi}(\Delta) = 0$, plugging in $1/\lambda$ in place of λ , and specializing to the ν -grading, the previous expression becomes:

$$(-1)^{d-1} F(k[\Delta], \nu) = \sum_{S \subseteq [d-1]} f'_S \prod_{i \in S} \frac{1}{\nu_i - 1}$$

Putting the above over the common denominator of $\prod_{i \in [d-1]} (\nu_i - 1)$ and multiplying by $(-1)^{d-1}$ gives us:

$$F(k[\Delta], \nu) = \sum_{S \subseteq [d-1]} \frac{f'_S \prod_{i \notin S} (\nu_i - 1)}{\prod_{i \in [d-1]} (1 - \nu_i)}$$

Comparing this with Equation (1) and noting that the denominators are equal, we have

$$\sum_{S \subseteq [d-1]} f'_S \prod_{i \notin S} (\nu_i - 1) = \sum_{S \subseteq [d-1]} h_S \prod_{i \in S} \nu_i = \sum_{S \subseteq [d-1]} h_{[d-1]-S} \prod_{i \notin S} \nu_i$$

which proves the result. \square

Now let Σ be a $(d-1)$ -dimensional shellable complex with face poset P_Σ and shelling order F_1, F_2, \dots, F_t , and for each i let P_i be the face poset of F_i . Let $A = [d-1]$, and set $\Delta = \Delta((P_\Sigma)_A)$. Note that Δ is simply the order complex of P_Σ once we remove the elements corresponding to the facets of Σ . Let $\Sigma_1, \Sigma_2, \dots, \Sigma_t$ be the c.e.d. of Δ given by Theorem 5.1.

Lemma 6.4. *Fix some Σ_i , and let λ_i be the labeling of P_i constructed in the proof of Theorem 5.1. Let $S, T \subseteq [d-1]$, and suppose S dominates T . Then there are at least as many maximal chains in Σ_i whose λ_i -labels have descent set S as chains whose λ_i -labels have descent set T .*

Proof. Let \mathbf{c} be a maximal chain in Σ_i with $\sigma = \lambda_i(\mathbf{c})$, and let $\sigma(j) < \sigma(j+1)$ be an ascent of its λ_i -label. Let \mathbf{c}' be the unique maximal chain of P_i that coincides with \mathbf{c} at every rank but j , and let x be the element of \mathbf{c}' of rank j . Then \mathbf{c}' must be a chain in Σ_i , since otherwise $\mathbf{c} = \Upsilon_{\lambda_i}(\mathbf{c}' - x)$ would not be a chain in Σ_i . So if τ is a permutation preceded by σ in the weak order, there is some maximal chain in Σ_i with τ as its λ_i -label. Now suppose σ has descent set T . If ϕ is as in Definition 2.10, it follows that Σ_i contains a chain whose λ_i -label is $\phi(\sigma)$. \square

Theorem 6.5. *Let Σ and Δ be as above, let $S, T \subseteq [d-1]$, and suppose that S dominates T . Then the flag h -vector of Δ satisfies $h_T \leq h_S$.*

Proof. First, note that $h_T(\Sigma_1) \leq h_S(\Sigma_1)$, since the poset associated to Σ_1 is just the Boolean lattice B_d . Now let $\Omega = \Sigma_1 \cup \Sigma_2 \cup \dots \cup \Sigma_{k-1}$ and suppose the result holds for Ω . Let $\Sigma'_k = \Sigma_k - \partial \Sigma_k - \emptyset$. Because Σ_k triangulates a ball, we can now use our earlier expression for the flag h -vector of a ball and invoke an argument similar to Chari's in [3]:

$$\begin{aligned} \sum_{S \subseteq [d-1]} h_S(\Omega \cup \Sigma_k) \prod_{i \notin S} \nu_i &= \sum_{S \subseteq [d-1]} f_S(\Omega \cup \Sigma_k) \prod_{i \notin S} (\nu_i - 1) \\ &= \sum_{S \subseteq [d-1]} f_S(\Omega) \prod_{i \notin S} (\nu_i - 1) + \sum_{S \subseteq [d-1]} f_S(\Sigma'_k) \prod_{i \notin S} (\nu_i - 1) \\ &= \sum_{S \subseteq [d-1]} h_S(\Omega) \prod_{i \notin S} \nu_i + \sum_{S \subseteq [d-1]} h_{[d-1]-S}(\Sigma_k) \prod_{i \notin S} \nu_i \quad (\text{by Lemma 6.3}) \\ &= \sum_{S \subseteq [d-1]} (h_S(\Omega) + h_{[d-1]-S}(\Sigma_k)) \prod_{i \notin S} \nu_i \end{aligned}$$

Thus, for all subsets $S \subseteq [d - 1]$,

$$h_S(\Omega \cup \Sigma_k) = h_S(\Omega) + h_{[d-1]-S}(\Sigma_k)$$

Because reverse lexicographic order of the maximal chains of Σ_k is a shelling (Theorem 3.1), $h_S(\Sigma_k)$ counts the number of maximal chains of Σ_k whose labels have *ascent* set S , and so $h_{[d-1]-S}(\Sigma_k)$ counts the number of maximal chains in P_k with descent set S . The result now follows from Lemma 6.4. \square

Proof of Theorem 6.2. Because P is a face poset, a linear inequality of its flag h-vector translates to a linear inequality of the h-vector of K . Since every h-vector of a Cohen-Macaulay complex is the h-vector of some shellable complex (see [13]), the result follows. \square

We now show that Theorem 6.2 cannot be extended to include posets whose order complexes are Cohen-Macaulay (or 2-CM, for that matter). First recall that a graded poset P is *Eulerian* if its Möbius function satisfies $\mu(x, y) = (-1)^{\text{rank}(y) - \text{rank}(x)}$ for all $x < y$. An Eulerian poset whose order complex is Cohen-Macaulay is called *Gorenstein**. It can be shown that the order complex of a Gorenstein* poset is 2-Cohen-Macaulay. For $S \subseteq [d - 1]$, define $w(S)$ to be the set of all $i \in [n]$ such that exactly one of i and $i + 1$ is in S . For instance, if $S = \{2, 3\} \subseteq [4]$ then $w(S) = \{1, 3\}$. Since Conjecture 2.3 from [12] was proven by Karu in [8], we can rephrase Proposition 2.8 from [12] as:

Proposition 6.6 ([12]). *If $S, T \subseteq [d - 1]$ are such that $h_T(\Delta) \leq h_S(\Delta)$ whenever Δ is the order complex of a Gorenstein* poset, then $w(T) \subseteq w(S)$.*

Now consider $S, T \subseteq [4]$ given by $S = \{1, 2\}$ and $T = \{1\}$. In [9], it is shown that S dominates T . However, $w(S) = \{2\}$ and $w(T) = \{1\}$, so $w(T) \not\subseteq w(S)$ and it is clear that we cannot extend Theorem 6.2 to include the wider class of Cohen-Macaulay posets (or even 2-CM posets).

Acknowledgements: This paper, which constituted part of the author's Ph.D. thesis, could not have been written without the encouragement, input, and patience of Ed Swartz. Thanks also to the anonymous referee for numerous helpful suggestions.

References

- [1] Anders Björner. The homology and shellability of matroids and geometric lattices. In *Matroid applications*, volume 40 of *Encyclopedia Math. Appl.*, pages 226–283. Cambridge Univ. Press, Cambridge, 1992.
- [2] Anders Björner and Michelle Wachs. On lexicographically shellable posets. *Trans. Amer. Math. Soc.*, 277(1):323–341, 1983.
- [3] Manoj K. Chari. Two decompositions in topological combinatorics with applications to matroid complexes. *Trans. Amer. Math. Soc.*, 349(10):3925–3943, 1997.

- [4] Gopal Danaraj and Victor Klee. Shellings of spheres and polytopes. *Duke Math. J.*, 41:443–451, 1974.
- [5] Timothy DeVries. The weak order and flag h -vector inequalities. Cornell University senior thesis, 2005.
- [6] Patricia Hersh and Ed Swartz. Coloring complexes and arrangements. *J. Algebraic Combin.*, 27(2):205–214, 2008.
- [7] Takayuki Hibi. Level rings and algebras with straightening laws. *J. Algebra*, 117(2):343–362, 1988.
- [8] Kalle Karu. The cd -index of fans and posets. *Compos. Math.*, 142(3):701–718, 2006.
- [9] Kathryn Nyman and Ed Swartz. Inequalities for the h -vectors and flag h -vectors of geometric lattices. *Discrete Comput. Geom.*, 32(4):533–548, 2004.
- [10] James G. Oxley. *Matroid theory*. Oxford Science Publications. The Clarendon Press Oxford University Press, New York, 1992.
- [11] Jay Schweig. A convex-ear decomposition for rank-selected subposets of supersolvable lattices. *SIAM J. Discrete Math.*, 23(2):1009–1022, 2009.
- [12] Richard P. Stanley. A survey of Eulerian posets. In *Polytopes: abstract, convex and computational (Scarborough, ON, 1993)*, volume 440 of *NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci.*, pages 301–333. Kluwer Acad. Publ., Dordrecht, 1994.
- [13] Richard P. Stanley. *Combinatorics and commutative algebra*, volume 41 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA, second edition, 1996.
- [14] Richard P. Stanley. *Enumerative combinatorics. Vol. 1*, volume 49 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1997. With a foreword by Gian-Carlo Rota, Corrected reprint of the 1986 original.
- [15] Ed Swartz. g -elements, finite buildings and higher Cohen-Macaulay connectivity. *J. Combin. Theory Ser. A*, 113(7):1305–1320, 2006.
- [16] Russ Woodroffe. Cubical convex ear decompositions. *Electron. J. Combin.*, 16(2, Special volume in honor of Anders Björner):Research Paper 17, 33, 2009.
- [17] Günter M. Ziegler. *Lectures on polytopes*, volume 152 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995.