

Math 2163, Practice Exam I

- Find $|a|$, $3a + 4b$, $a \cdot b$, $a \times b$ where
 - $a = \langle -3, 1, 2 \rangle$, $b = \langle 6, 1, 7 \rangle$;
 - $a = i - 2j + k$, $b = j + 2k$.
- Find a unit vector that has the same direction as $12i - 5j$.
- Find the angle between vectors a and b , determine whether they are orthogonal, parallel, or skew:
 - $a = \langle 6, -3, 2 \rangle$, $b = \langle 2, 1, -2 \rangle$;
 - $a = 2i - j + k$, $b = 3i + 2j - k$;
 - $a = \langle 4, 6 \rangle$, $b = \langle -3, 2 \rangle$;
 - $a = 2i + 6j - 4k$, $b = -3i - 9j + 6k$;
 - $a = \langle a, b, c \rangle$, $b = \langle -b, a, 0 \rangle$.
- Find the direction cosines of the vector $\langle 1, -2, -1 \rangle$.
- Find the scalar and vector projection of $\langle -1, -2, 2 \rangle$ onto $\langle 3, 3, 4 \rangle$.
- Find the two unit vectors orthogonal to both $i + j + k$ and $2i + k$.
- Find the volume of the parallelepiped decided by $\langle 1, 1, -1 \rangle$, $\langle 1, -1, 1 \rangle$ and $\langle -1, 1, 1 \rangle$.
- Find the lines specified by:
 - passes through the point $(1, 0, -3)$ and parallel to the vector $\langle 2, -4, 5 \rangle$;
 - passes through the point $(-2, 4, 1)$ and perpendicular to the plane $x + y - z = 5$;
 - passes through two points $(1, 1, 3)$ and $(2, -1, 0)$;
 - passes through the point $(2, 1, 0)$ and perpendicular to both $i + j + k$ and $i - j$;
 - the line of intersection of the planes $x + y + z = 1$ and $x + z = 0$.
- Find the plane specified by:
 - passes through the point $(4, 1, 2)$ and perpendicular to the vector $i + j + 2k$;
 - passes through the point $(4, 1, 2)$ and parallel to the plane $x - y + z = 5$;
 - passes through three points $(3, -1, 2)$, $(8, 2, 4)$ and $(-1, -2, -3)$;
 - passes through a point $(1, 3, 0)$ and contains the line $x = 1 + t$, $y = 2t$, $z = -1 - t$;
- Find the domain of the following functions:
 - $f(x, y) = \sqrt{y - x} \ln(y + x)$;

(b) $f(x, y) = \sqrt{16 - x^2 - 16y^2}$;

(c) $f(x, y) = \frac{x-3y}{x^2-y}$

11. Show that the following limits does not exist:

(a) $\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 + 2y^2}{2x^2 + y^2}$;

(b) $\lim_{(x,y) \rightarrow (0,0)} \frac{x^3y}{x^4 + y^4}$;

(c) $\lim_{(x,y) \rightarrow (0,0)} \frac{\sin(xy)}{x^2 - y^2}$

12. Find all first order partial derivatives for:

(a) $f(x, y) = x^y$;

(b) $f(x, y) = \frac{xy^2}{x^2+y^2}$;

(c) $f(x, y) = \int_x^y \cos t \, dt$;

(d) $f(x, y, z) = x^2e^{yz}$;

(e) $f(x, y, z) = x \tan(xy)$;

(f) $z = \sin u$ and $u = y/x$;

(g) $z = xy + y^2$ and $x = 2t, y = 1 - t$;

(h) $z = xy + yz + zx$ and $x = st, y = e^{st}, z = t^2$;

(i) $xy^2 + yz^2 + zx^2 = 3$ and z is a function of x and y ;

(j) $xyz = \cos(xyz)$ and z is a function of x and y ;

13. Find the following higher order derivatives:

(a) $z = x \sin y$, find $\frac{\partial^2 z}{\partial y^2}$;

(b) $f(x, y) = e^{xy^2}$, find f_{yy} ;

(c) $f(r, s, t) = r \ln(rs^2t^3)$, find f_{rss} and f_{rst} .

14. Find the total differential dz of $z = e^s \sin t$.

15. Given $z = \ln(2x + y)$, find the tangent plane at $(-1, 3, 0)$.

16. Find the linear approximation of $f(x, y) = xe^{xy}$ at $(1, 0)$ and use it to approximate $f(1.1, -0.1)$.