

MATH 4063-5023
Homework Set 5

1. Suppose that V is a finitely generated vector space and $\phi : V \rightarrow W$ is a linear transformation. Show that $\text{im}(\phi) \subset W$ is finitely generated.

- Since V is finitely generated it has a finite basis; say, $B = \{v_1, \dots, v_n\}$ is a basis for V . Now

$$\begin{aligned}\text{im}(\phi) &\equiv \{w \in W \mid w = \phi(v) \quad , \quad v \in V\} \\ &= \{w \in W \mid w = \phi(\alpha_1 v_1 + \dots + \alpha_n v_n) \quad , \quad \alpha_i \in \mathbb{F}\}\end{aligned}$$

since each vector in V can be expressed as a linear combination of the $v_i \in B$. But then

$$\begin{aligned}\text{im}(\phi) &= \{w \in W \mid w = \alpha_1 \phi(v_1) + \dots + \alpha_n \phi(v_n) \quad , \quad \alpha_i \in \mathbb{F}\} \quad \text{since } \phi \text{ is a linear transformation} \\ &= \text{span}(\phi(v_1), \dots, \phi(v_n))\end{aligned}$$

So $\text{im}(\phi)$ is generated by $\phi(v_1), \dots, \phi(v_n)$. □

2. Suppose S is a subspace of a finitely generated vector space V . Show that V/S is finitely generated.

- Let $p_S : V \rightarrow V/S$ be the canonical projection. This is a linear transformation between V and V/S with image V/S . Therefore, by Problem 1 above, since V is finitely generated, the image V/S of p_S must be finitely generated. □

3. Suppose S is a subspace of a finitely generated vector space V , find a basis for V/S .

- Since S is a subspace of a f.g. vector space, S has a basis $B_S = \{b_1, \dots, b_k\}$. Moreover, the basis for the subspace S can always be extended to a basis for V . Let

$$B_V = \{b_1, \dots, b_k, b'_{k+1}, \dots, b'_n\}$$

be such a basis for V . From Problems (1) and (2) above, we know that

$$V/S = \text{im}(p_S) = \text{span}(p_S(b_1), \dots, p_S(b_k), p_S(b'_{k+1}), \dots, p_S(b'_n))$$

Now the for $1 \leq i \leq k$, $b_i \in S = \ker(p_S)$ and so

$$V/S = \text{span}(\mathbf{0}_{V/S}, \dots, \mathbf{0}_{V/S}, p_S(b'_{k+1}), \dots, p_S(b'_n)) = \text{span}(p_S(b'_{k+1}), \dots, p_S(b'_n))$$

So the vectors $p_S(b'_{k+1}), \dots, p_S(b'_n)$ span V/S . I claim they are also linearly independent. Indeed,

$$\begin{aligned}\mathbf{0}_{V/S} &= \alpha_{k+1} p_S(b'_{k+1}) + \dots + \alpha_n p_S(b'_n) \\ &= p_S(\alpha_{k+1} b'_{k+1} + \dots + \alpha_n b'_n) \\ \Rightarrow \quad \alpha_{k+1} b'_{k+1} + \dots + \alpha_n b'_n &\in \ker(p_S) = S\end{aligned}$$

But a vector $\alpha_1 b_1 + \dots + \alpha_k b_k + \alpha_{k+1} b'_{k+1} + \dots + \alpha_n b'_n$ can be in S only if $0 = \alpha_{k+1}$, $0 = \alpha_{k+2}$, \dots , $0 = \alpha_n$. Hence,

$$\mathbf{0}_{V/S} = \alpha_{k+1} p_S(b'_{k+1}) + \dots + \alpha_n p_S(b'_n) \Rightarrow 0 = \alpha_{k+1} = \alpha_{k+2} = \dots = \alpha_n$$

hence the vectors $p_S(b'_{k+1}), \dots, p_S(b'_n)$ are linearly independent. □

4. Suppose S is a subspace of a finitely generated vector space V , show that $\dim(V) = \dim(S) + \dim(V/S)$.

- From the theory of linear transformations we have, since the canonical projection $p_S : V \rightarrow V/S$ is a linear transformation

$$\dim(V) = \dim(\text{im}(p_S)) + \dim(\ker(p_S))$$

On the other hand,

$$\begin{aligned}\ker(p_S) &= S \\ \text{im}(p_S) &= V/S\end{aligned}$$

and so

$$\dim(V) = \dim(V/S) + \dim S \quad .$$

□

5. Suppose $\phi : V \rightarrow W$ is a linear transformation between two finite-dimensional vector spaces. Show that $\text{im}(\phi)$ is isomorphic to $V/\ker(\phi)$. (Hint: two finite-dimensional vector spaces are isomorphic if and only if they have the same dimension.)

- Regarding $\ker \phi$ as a subspace of V we have a canonical projection $p_{\ker \phi} : V \rightarrow V/\ker \phi$, and from Problem 4 above we have

$$(i) \quad \dim V = \dim(V/\ker \phi) + \dim(\ker \phi)$$

On the other hand, from our general theory of linear transformations

$$(ii) \quad \dim(V) = \dim(\text{im}(\phi)) + \dim(\ker \phi)$$

Comparing (i) with (ii) we conclude

$$\dim(V/\ker \phi) = \dim(\text{im}(\phi))$$

Because $V/\ker \phi$ and $\text{im}(\phi)$ are finite-dimensional and share the same dimension, they must be isomorphic. □