

ON THE HEIGHTS OF TOTALLY p -ADIC NUMBERS

PAUL FILI

ABSTRACT. Bombieri and Zannier established lower and upper bounds for the limit infimum of the Weil height in fields of totally p -adic numbers and generalizations thereof. In this paper, we use potential theoretic techniques to generalize the upper bounds from their paper and, under the assumption of integrality, to improve slightly upon their bounds.

1. STATEMENT OF RESULTS

Recall that an algebraic number is said to be *totally p -adic* if its image lies in \mathbb{Q}_p for any embedding $\mathbb{Q} \hookrightarrow \mathbb{C}_p$, where \mathbb{C}_p denotes the completion of an algebraic closure of \mathbb{Q}_p . This is analogous to the usual definition of a totally real number, however, unlike \mathbb{C}/\mathbb{R} , the extension $\mathbb{C}_p/\mathbb{Q}_p$ is of infinite degree, so in fact we can make an even broader generalization:

Definition 1. Let L_p/\mathbb{Q}_p be a (finite) Galois extension for $p \leq \infty$ a rational prime. We say $\alpha \in \overline{\mathbb{Q}}$ is *totally L_p* if all Galois conjugates of α lie in $L_p \subset \mathbb{C}_p$.

More generally, when our objects are defined over an arbitrary number field K , we make the following definition:

Definition 2. Fix a base number field K , and let S be a set of places of K . For each $v \in S$, we choose a Galois extension L_v/K_v . We say that α is *totally L_S/K* if, for each $v \in S$, all of the K -Galois conjugates of α lie in L_v .

Notice that α is totally L_S/K if and only if the minimal polynomial for α over K splits in L_v for each $v \in S$. In our terminology, a number being totally real is equivalent to being totally \mathbb{R} (or totally \mathbb{R}/\mathbb{Q}), and being totally p -adic is equivalent to being totally \mathbb{Q}_p (totally \mathbb{Q}_p/\mathbb{Q}). Notice that the set of all totally L_S/K numbers form a normal extension of K (typically of infinite degree).

Bombieri and Zannier [3] studied the question of what the limit infimum of the Weil height was in these fields when the base field was assumed to be \mathbb{Q} , and they proved the following:

Theorem (Bombieri and Zannier 2001). *Let S be the set of finite rational primes and for each $p \in S$, let L_p/\mathbb{Q}_p be a chosen Galois extension. Suppose L/\mathbb{Q} is a normal extension (possibly of infinite degree) such that L embeds in L_p for each $p \in S$. Then*

$$\liminf_{\alpha \in L} h(\alpha) \geq \frac{1}{2} \sum_{p \in S} \frac{\log p}{e_p(p^{f_p} + 1)}$$

Date: May 8, 2013.

2010 Mathematics Subject Classification. 11G50, 11R06, 37P30.

Key words and phrases. Weil height, totally p -adic, potential theory, Fekete-Szegő theorem.

where e_p and f_p denote the ramification and inertial degrees of L_p/\mathbb{Q}_p , respectively.

Further, in the case where $S = \{p_1, \dots, p_n\}$ is a finite set of nonarchimedean rational primes with $L_{p_i} = \mathbb{Q}_{p_i}$ for each i , if we let L be the field of all totally L_S/\mathbb{Q} numbers, then we have

$$\liminf_{\alpha \in L} h(\alpha) \leq \sum_{i=1}^n \frac{\log p_i}{p_i - 1}.$$

We note that in Bombieri and Zannier's result, by the assumption that our extensions are normal, it follows that every p -adic completion of L has ramification and inertial degree less than or equal to that of L_p . More simply, we might take L to be the field of all totally L_S/\mathbb{Q} numbers above and recover the same result.

Smyth [8, 9] and Flammang [4] proved analogous results in the totally real case for the limit infimum of the height, however, they imposed the additional hypothesis that one consider only totally real algebraic integers.

This note has two aims: first, to prove a slightly sharper lower bound by adding the hypothesis that we consider only algebraic integers, and second, to generalize the upper bound from Bombieri and Zannier's paper by the aid of the Fekete-Szegő theorem with splitting conditions, as formulated and proven by Rumely [5, 6]. Our results are the following:

Theorem 1. *Fix a number field K , a set of nonarchimedean places S of K , and a choice of Galois extension L_v/K_v for each $v \in S$. Let L be the field of all totally L_S/K numbers, and O_L denote its ring of integers. Then*

$$\liminf_{\alpha \in O_L} h(\alpha) \geq \frac{1}{2} \sum_{v \in S} N_v \cdot \frac{\log p_v}{e_v(q_v^{f_v} - 1)}$$

where $N_v = [K_v : \mathbb{Q}_v]/[K : \mathbb{Q}]$, e_v and f_v denote the ramification and inertial degrees of L_v/K_v , respectively, and q_v denotes the order of the residue field of K_v , and p_v is rational prime above which v lies.

As Bombieri and Zannier remark in their note, if the above sum diverges (which may happen if we allow S to be infinite), then it follows that the set of totally L_S/K algebraic integers satisfies the Northcott property (there are only a finite number with height below any fixed constant).

If we restrict our attention to extensions normal over \mathbb{Q} , as Bombieri and Zannier do, our result reduces to:

Corollary 1. *Fix a set of finite rational primes S and a choice of Galois extension L_p/\mathbb{Q}_p for each $p \in S$. Let L be the field of all totally L_S numbers, and O_L denote its ring of integers. Then*

$$\liminf_{\alpha \in O_L} h(\alpha) \geq \frac{1}{2} \sum_{p \in S} \frac{\log p}{e_p(p^{f_p} - 1)}$$

where e_p and f_p denote the ramification and inertial degrees of L_p/\mathbb{Q}_p , respectively.

Theorem 2. *Fix a number field K , a finite set of nonarchimedean places S of K , and a choice of Galois extension L_v/K_v for each $v \in S$ with ramification degree e_v , inertial degree f_v , and residue field degree q_v and characteristic p_v . Let L be the field of all totally L_S/K numbers. Then*

$$\liminf_{\alpha \in L} h(\alpha) \leq \sum_{v \in S} N_v \cdot \frac{\log p_v}{e_v(q_v^{f_v} - 1)}.$$

where $N_v = [K_v : \mathbb{Q}_v]/[K : \mathbb{Q}]$.

Notice that this differs from our lower bound only by the factor of $1/2$.

One interesting thing to note in Theorem 1 is that the shape of our bound comes directly from the formula for the logarithmic capacity of the ring of integers of L_v at each place. This connection to potential theory gives an interesting explanation for the shape of the observed bound, and suggests that by determining minimal energy measures on L_v one might further improve the bounds above. In fact, this connection inspires us to conjecture that in fact the upper bound in Theorem 2 is sharp:

Conjecture 1. *Fix a number field K , a finite set of nonarchimedean places S of K , and a choice of Galois extension L_v/K_v for each $v \in S$ with ramification degree e_v , inertial degree f_v , and residue field of degree q_v and characteristic p_v . Let L be the field of all totally L_S/K numbers and O_L its ring of integers. Then*

$$\liminf_{\alpha \in O_L} h(\alpha) = \sum_{v \in S} N_v \cdot \frac{\log p_v}{e_v(q_v^{f_v} - 1)}.$$

where $N_v = [K_v : \mathbb{Q}_v]/[K : \mathbb{Q}]$.

It is still an interesting open question whether the limit infimum for all numbers can be in fact achieved with algebraic integers or not. In this direction, we will note only that the result of Bombieri and Zannier, when read as an equidistribution result, seems to indicate that totally p -adic points of low height should be distributing evenly in the residue classes of $\mathbb{P}^1(\mathbb{F}_p)$. This seems to suggest that the limit infimum over an entire field L of the type constructed above might be smaller than that obtained by algebraic integers; however, all of the smallest known limit points, both for the height of totally real numbers (see [8]) and the height of totally p -adic numbers (as in Theorem 2) are in fact achieved by sequences of algebraic integers.

2. PROOFS

Proof of Theorem 1. We now proceed to prove Theorem 1. Recall that we have fixed a base number field K , and let M_K denotes the places of K . First, let us note that if S is infinite, we can take a limit over increasing finite subsets of S , and the general result will follow, so we may as well assume that S is finite in our proof. For convenience let

$$N_v = \frac{[K_v : \mathbb{Q}_v]}{[K : \mathbb{Q}]}.$$

We recall M. Baker's reformulation of Mahler's inequality from [1], namely, if α is an algebraic number and $\alpha_1, \dots, \alpha_n$ denote its K -Galois conjugates, then

$$(1) \quad \frac{1}{n(n-1)} \sum_{i \neq j} g_v(x, y) \geq \begin{cases} -N_v \cdot \frac{\log n}{n-1} & \text{if } v \mid \infty, \\ 0 & \text{otherwise} \end{cases}$$

where

$$g_p(x, y) = \log^+ |x|_v + \log^+ |y|_v - \log^+ |x - y|_v,$$

and our absolute values are normalized so that $|\cdot|_v = \|\cdot\|_v^{N_v}$ where $\|\cdot\|_v$ extends the usual absolute of \mathbb{Q} over which it lies (thus the absolute values $|\cdot|_v$ satisfy the product formula, and $h(\alpha) = \sum_v \log^+ |\alpha|_v$). We assume in our theorem that α is

integral. The key role is played by the discriminant here. At the place $v \in M_K$ (p here being the rational prime over which the place v lies), all of the conjugates of α lie in ring of integers O_{L_v} . By [7, Example 4.1.24], we have that the v -adic logarithmic capacity of O_{L_v} with respect to the point ∞ , which we will denote $\gamma_{\infty,v}(O_{L_v})$, satisfies

$$\log \gamma_{\infty,v}(O_{L_v}) = -N_v \cdot \frac{\log p_v}{e_v(q_v^{f_v} - 1)} < 0,$$

where q_v denotes the order of the residue field of O_{L_v} and p_v its characteristic.¹ We can also view this as the transfinite diameter of O_{L_v} ; more specifically, if we adopt the notation of [2, Ch. 6] and let

$$\log d_n(O_{L_v})_{\infty} = \sup_{z_1, \dots, z_n \in O_{L_v}} \frac{1}{n(n-1)} \log |z_i - z_j|_v,$$

then by [2, Lemma 6.21 and Theorem 6.23] it follows that²

$$\lim_{n \rightarrow \infty} \log d_n(O_{L_v})_{\infty} = \log \gamma_{\infty,v}(O_{L_v}).$$

Let $\{\alpha^{(k)}\}_{k=1}^{\infty}$ denote a sequence of algebraic numbers such that

$$\lim_{k \rightarrow \infty} h(\alpha^{(k)}) = \liminf_{\alpha \in O_L} h(\alpha),$$

and let n_k denote the number of K -Galois conjugates of $\alpha^{(k)}$, which we denote $\alpha_1^{(k)}, \dots, \alpha_{n_k}^{(k)}$. Since the $\alpha^{(k)}$ have bounded height, it follows from Northcott's theorem that $n_k \rightarrow \infty$ as $k \rightarrow \infty$. By definition of $d_n(O_{L_v})_{\infty}$, we have

$$\frac{1}{n_k(n_k - 1)} \sum_{i \neq j} \log |\alpha_i^{(k)} - \alpha_j^{(k)}|_v \leq \log d_{n_k}(O_{L_v})_{\infty}$$

for any totally L_S/K integer and $v \in S$, so it follows that

$$\limsup_{k \rightarrow \infty} \sum_{v \in S} \frac{1}{n_k(n_k - 1)} \sum_{i \neq j} \log |\alpha_i^{(k)} - \alpha_j^{(k)}|_v \leq \sum_{v \in S} \log \gamma_{\infty,v}(O_{L_v}).$$

Therefore, by the product formula, using the assumption of integrality,

$$\liminf_{k \rightarrow \infty} \sum_{\substack{w \in M_K \\ w | \infty}} \frac{1}{n_k(n_k - 1)} \sum_{i \neq j} \log |\alpha_i^{(k)} - \alpha_j^{(k)}|_w \geq - \sum_{v \in S} \log \gamma_{\infty,v}(O_{L_v}).$$

It now follows from (1) that

$$\begin{aligned} \liminf_{k \rightarrow \infty} 2h(\alpha^{(k)}) &\geq \liminf_{k \rightarrow \infty} \sum_{w | \infty} \log^+ |\alpha^{(k)}|_w \\ &\geq - \sum_{v \in S} \log \gamma_{\infty,v}(O_{L_v}) = \sum_{v \in S} N_v \cdot \frac{\log p_v}{e_v(q_v^{f_v} - 1)}. \end{aligned}$$

¹In the text [7] the v -adic absolute value $|\cdot|_v$ which is used to compute the capacity is normalized to agree with the modulus with respect to the additive Haar measure, so in the text a $\log q_v$ appears in the numerator instead of $\log p_v$. We also wish to draw the reader's attention to the fact that the ring of integers O_{L_v} is 'strictly smaller' than the closed unit disc in $\mathbb{P}^1(\mathbb{C}_v)$, which has logarithmic capacity $\log \gamma_{\infty,v}(\mathcal{D}_v(0,1)) = 0$.

²Note that here for convenience our absolute values are normalized so that we have the same normalization factors as in computing the absolute logarithmic Weil height.

Upon dividing each side by 2, the result follows. \square

Proof of Theorem 2. We will prove the result by applying the adelic Fekete-Szegő theorem with splitting conditions [6, Theorem 2.1] (see also [5] as well as [2, Theorem 6.27] for an interpretation in terms of the Berkovich topology). For each $v \in S$, let $O_{L_v} \subset L_v$ denote the ring of integers in L_v . Notice that $O_{L_v} \subset \mathbb{P}^1(\mathbb{C}_p)$ is a compact set and let $\mathcal{D}_v(0, 1) \subset \mathbb{P}^1(\mathbb{C}_v)$ be the usual Berkovich unit disc. Let $\mathbb{E} \subset \prod_{v \in M_K} \mathbb{A}^1(\mathbb{C}_v)$ be the adelic Berkovich set given by

$$\mathbb{E} = \prod_{v \in M_K} E_v \quad \text{where} \quad E_v = \begin{cases} O_{L_v} & \text{for } v \in S \\ \mathcal{D}_v(0, 1) & \text{for } v \notin S, \\ \mathcal{D}\left(0, \prod_{v \in S} q_v^{N_v/e_v(q_v^{f_v}-1)}\right) & \text{for } v \mid \infty. \end{cases}$$

By [7, Example 4.1.24] we have normalized v -adic logarithmic capacity

$$\log \gamma_{\infty, v}(O_{L_v}) = -N_v \cdot \frac{\log p_v}{e_v(q_v^{f_v} - 1)} \quad \text{for each } v \in S,$$

where γ_{∞} denotes the capacity relative to $\mathcal{X} = \{\infty\}$ in the notation of [6]. At the infinite places, we have

$$\sum_{w \mid \infty} \log \gamma_{\infty, w}(\mathcal{D}(0, \prod_{v \in S} q_v^{N_v/e_v(q_v^{f_v}-1)})) = \sum_{v \in S} N_v \cdot \frac{\log p_v}{e_v(q_v^{f_v} - 1)},$$

so that the adelic capacity, that is, the product of all of the normalized capacities, is

$$\gamma_{\infty}(\mathbb{E}) = \prod_{v \in M_K} \gamma_{\infty, v}(E_v) = 1.$$

Further, by our assumptions, \mathbb{E} is stable under the action of the continuous Galois automorphisms of \mathbb{C}_v/K_v at each place. For $\epsilon > 0$, we consider the Berkovich adelic neighborhood

$$\mathbb{U} = \prod_{w \in M_K} U_w \quad \text{where} \quad U_w = \begin{cases} E_w & \text{for } w \nmid \infty \\ \mathcal{D}\left(0, e^{\epsilon} \prod_{v \in S} q_v^{N_v/e_v(q_v^{f_v}-1)}\right)^- & \text{for } w \mid \infty, \end{cases}$$

where $\mathcal{D}(a, r)^-$ denotes the usual open disc centered at a of radius r . Then we can apply the adelic Fekete-Szegő theorem with splitting conditions as formulated in [6, Theorem 2.1], there are infinitely many algebraic numbers in \overline{K} all of whose conjugates lie in \mathbb{U} with the additional condition that all \mathbb{C}_v/K_v conjugates lie in $U_v = E_v = O_{L_v}$ for each $v \in S$. Clearly, the only contribution to the height of such numbers comes from the archimedean places, and as a result, we have

$$h(\alpha) \leq \sum_{v \in S} N_v \cdot \frac{\log p_v}{e_v(q_v^{f_v} - 1)} + \epsilon.$$

Thus by letting $\epsilon \rightarrow 0$ we can generate a sequence of numbers (in fact, algebraic integers) with the limit infimum of the height bounded by the desired constant. \square

REFERENCES

- [1] M. Baker. A lower bound for average values of dynamical Green's functions. *Math. Res. Lett.*, 13(2-3):245–257, 2006.
- [2] M. Baker and R. Rumely. *Potential theory and dynamics on the Berkovich projective line*, volume 159 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2010.
- [3] E. Bombieri and U. Zannier. A note on heights in certain infinite extensions of \mathbb{Q} . *Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl.*, 12:5–14, 2001.
- [4] V. Flammang. Two new points in the spectrum of the absolute Mahler measure of totally positive algebraic integers. *Math. Comp.*, 65(213):307–311, 1996.
- [5] R. Rumely. The Fekete-Szegő theorem with splitting conditions. I. *Acta Arith.*, 93(2):99–116, 2000.
- [6] R. Rumely. The Fekete-Szegő theorem with splitting conditions. II. *Acta Arith.*, 103(4):347–410, 2002.
- [7] R. S. Rumely. *Capacity theory on algebraic curves*, volume 1378 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1989.
- [8] C. J. Smyth. On the measure of totally real algebraic integers. *J. Austral. Math. Soc. Ser. A*, 30(2):137–149, 1980/81.
- [9] C. J. Smyth. On the measure of totally real algebraic integers. II. *Math. Comp.*, 37(155):205–208, 1981.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ROCHESTER, ROCHESTER, NY 14627
E-mail address: `fili@post.harvard.edu`