

LCM LATTICES SUPPORTING PURE RESOLUTIONS

CHRISTOPHER A. FRANCISCO, JEFFREY MERMIN, AND JAY SCHWEIG

ABSTRACT. We characterize the lcm lattices that support a monomial ideal with a pure resolution. Given such a lattice, we provide a construction that yields a monomial ideal with that lcm lattice and whose minimal free resolution is pure.

1. INTRODUCTION

Understanding the minimal free resolutions of monomial ideals has been a deceptively difficult problem that has occupied researchers for over 50 years. In recent decades, combinatorial techniques have both helped shed new light on methods for constructing minimal resolutions and illustrated some of the problems that arise. For example, while simplicial and cellular resolutions have proven useful in many situations, Velasco showed that there are monomial ideals whose minimal resolutions are not even supported on a CW-complex [V].

Nevertheless, combinatorial and topological methods still allow one to understand the minimal resolutions of monomial ideals at the cost of computing homologies. For example, formulas of Hochster [H] and Bayer, Charalambous, and Popescu [BCP] yield multigraded Betti numbers for any monomial ideal provided that one can compute the relevant homology groups for certain simplicial complexes. Gasharov, Peeva, and Welker introduce the lcm lattice in the context of resolutions in [GPW]. The combinatorial structure of the lcm lattice of an ideal determines its total Betti numbers and projective dimension. (See [Pe, Section 58] for background on lcm lattices.)

However, two ideals with isomorphic lcm lattices need not have the same graded Betti numbers. One particularly interesting type of free resolution is the class of pure resolutions, those for which at each step in the resolution, there is only a single shift. Pure resolutions play a vital role in Boij-Söderberg theory, which decomposes free resolutions in an especially nice way. In this paper, we ask:

Question 1.1. When is a lattice the lcm lattice of a monomial ideal with a pure resolution?

We provide a complete answer to this question in Theorem 3.8, using a topological condition on the lattice that we call *homologically monotonic* (HM).

Theorem 1.2 (Theorem 3.8). *Let Λ be an atomic lattice. Then there exists an ideal I with pure resolution such that $\text{lcm}(I) = \Lambda$ if and only if Λ is HM.*

As we note in Remark 3.2, geometric and distributive (or, more generally, supersolvable) lattices are HM, and thus Theorem 3.8 encompasses a wide range of lattices. Using the proof

of Theorem 3.8, given any lattice Λ that is HM, we explain how to construct a monomial ideal with a pure resolution and lcm lattice Λ .

In Section 2, which follows the work of Phan [Ph] and Mapes [M], we review the necessary background on lcm lattices and describe the labeling techniques we will use in our proofs. The main results are in Section 3.

2. BACKGROUND AND NOTATION

Throughout the paper, let $R = \mathbb{k}[x_1, x_2, \dots, x_n]$ be the polynomial ring with sufficiently many variables over a field \mathbb{k} . Let Λ denote a finite atomic lattice, defined below.

Definition 2.1. A finite poset Λ is called a *finite atomic lattice* if it satisfies the following properties:

- (1) Every subset $S \subseteq \Lambda$ has a unique least upper bound $\bigvee(S)$, called the *join* of S , and a unique greatest lower bound $\bigwedge(S)$, called the *meet* of S .
- (2) Every element of Λ is a join of atoms. (Recall that an atom is an element that covers the least element $\hat{0}$.)

Note that, if Λ is a finite atomic lattice, there are unique maximal and minimal elements $\hat{1} = \bigvee(\Lambda)$ and $\hat{0} = \bigwedge(\Lambda)$.

Finally, an element $v \in \Lambda \setminus \{\hat{1}\}$ is called *meet-irreducible* if it cannot be written as the meet of two larger elements.

Let I be a monomial ideal with minimal generating set $G = (m_1, \dots, m_t)$. Put $\Lambda = \{\mu : \mu = \text{lcm}(S) \text{ for some } S \subseteq G\}$. Then Λ , ordered by divisibility, forms a finite atomic lattice. The generators of I are the atoms, and $\hat{0} = 1$. In this case, Λ is called the *lcm lattice* of I , denoted $\text{lcm}(I) = \Lambda$. See Figure 1 for an example.

Given a finite atomic lattice, there are two standard ways to associate monomials to some of its vertices (see [M] for details), and we are in the habit of referring to both associations as *labelings*. Unfortunately, the present paper will require these two labelings to interact with one another in complicated ways, so to avoid ambiguity we will avoid words like “labeling” in favor of “tags” for one association, and “multidegree” for the other.

Definition 2.2. A *multigraded lattice* is a pair (Λ, mdeg) , where Λ is a finite atomic lattice, and mdeg is a function from the vertices of Λ to the monomials of R which satisfies the properties below. (We refer to $\text{mdeg}(v)$ as the *multidegree* of v .)

- (1) Atoms of Λ have incomparable multidegree.
- (2) $\text{mdeg}(w_1 \vee w_2) = \text{lcm}(\text{mdeg}(w_1), \text{mdeg}(w_2))$.
- (3) $\text{mdeg}(\hat{0}) = 1$.
- (4) For all v , $\text{mdeg}(v) = \text{lcm}(\text{mdeg}(a) : a \leq v \text{ and } a \text{ is an atom})$.

Observe that (4) is equivalent to (2) and (3).

Example 2.3. The lcm lattice of the ideal $I = (c^2d, acde, a^2b^2, abce)$ is shown in Figure 1. The multigrading function assigns each vertex its own multidegree; the lcm lattice together with this natural multigrading is a multigraded lattice.




FIGURE 1. The (multigraded) lcm lattice of the ideal $I = (c^2d, acde, a^2b^2, abce)$.

Definition 2.4. A *tagged lattice* is a pair (Λ, τ) , where Λ is a finite atomic lattice and τ is a function from the vertices of Λ to the monomials of R which satisfies the properties below. (We refer to $\tau(v)$ as the *tag* of v .)

- (1) If v is meet-irreducible, then $\tau(v) \neq 1$.
- (2) For every variable x , the set $\{v : x \text{ divides } \tau(v)\}$ forms a chain in Λ .
- (3) $\tau(\hat{1}) = 1$.




FIGURE 2. A tagged lattice.

Example 2.5. The lattice in Figure 2 is tagged. The underlying lattice Λ is the same as in Figure 1. The tagging function τ assigns each unmarked vertex to the monomial 1, and each marked vertex to its marking. Observe that every meet-irreducible vertex has a non-trivial tag, and that, while the variables a , b , and c each appear in multiple tags, these tags always form a chain. We will see below that the tagged lattice of Figure 2 and the multigraded lattice of Figure 1 contain the same information.

There is a natural duality between tagged and multigraded lattices, given by the following constructions. (We use only Construction 2.6 in this paper but state Construction 2.7 for completeness.)

Construction 2.6 (Phan, Mapes). Suppose (Λ, τ) is a tagged lattice. For each vertex $v \in \Lambda$, set $\tau^m(v) = \prod_{w \not\leq v} \tau(w)$. Then the pair (Λ, τ^m) is a multigraded lattice.

Construction 2.7 (Mapes). Suppose (Λ, mdeg) is a multigraded lattice. For each vertex $v \in \Lambda$ and each variable x_i , set $\ell_i(v) = \min_{w \not\leq v} (\max(\ell : x_i^\ell \text{ divides } \text{mdeg}(w)))$ and $s_i(v) = \max(s : x_i^s \text{ divides } \text{mdeg}(v))$. Finally, set $\tau(v) = \prod x_i^{\ell_i(v) - s_i(v)}$. Then the pair (Λ, τ) is a tagged lattice.

Proposition 2.8 (Mapes). *Constructions 2.6 and 2.7 are inverse bijections. In particular, there is a one-to-one correspondence between monomial ideals, multigraded lattices, and tagged lattices.*

Proof. For the inverse bijection, see [M]. The correspondence to monomial ideals associates a monomial ideal to its (multigraded) lcm lattice, and a multigraded lattice to the ideal generated by the multidegrees of its atoms. \square

The lattices of Examples 2.3 and 2.5 are associated to one another via Constructions 2.6 and 2.7.




FIGURE 3. The lattice of the examples.

Example 2.9. Let $v \in \Lambda$ be as in Figure 3, where Λ is the underlying lattice of our running examples. In Example 2.5, the vertices greater than or equal to v have tags b and ab , so the multidegree of v is the product of the other tags, $\tau^m(v) = (a)(c)(e)(d)(c) = ac^2de$. In Example 2.3, the multidegree of v is ac^2de , which is not divisible by b . Since every vertex not less than or equal to v has multidegree divisible by b , it follows that the tag of v is divisible by b . Since there are incomparable elements with multidegrees divisible by the same power of every other variable, it follows that the tag of v is in fact equal to b . Analyzing every other vertex in this manner, we see that the lattices of Examples 2.3 and 2.5 correspond under Constructions 2.6 and 2.7.

Notation 2.10. Throughout the paper, we will study the (multigraded lattice and) monomial ideal coming from a tagged lattice via Construction 2.6. In order to avoid notational monstrosities like “ $\deg(\text{mdeg}(v))$,” we refer to the multigrading by τ^m . Thus our multigraded lattices are (Λ, τ^m) , and in Example 2.9 we have $\tau^m(v) = ac^2de$ and $\deg(\tau^m(v)) = 5$.

The main algebraic motivation for studying lattices is that the Betti numbers of a monomial ideal are encoded in the poset structure of its lcm lattice. For $x < y$ in a poset P , recall that the *order complex* of the open interval (x, y) is the simplicial complex whose faces are chains in P between x and y . The following result is normally stated for the quotient S/I rather than the ideal I ; see, e.g., [Pe, Theorem 58.8].

Theorem 2.11 (Hochster’s formula for lcm lattices). *Let I be a monomial ideal, and let (Λ, τ^m) be its multigraded lcm lattice. Fix $v \neq \hat{0} \in \Lambda$, with $\tau^m(v) = \mu$. Then the multigraded Betti numbers of I with multidegree μ are given by the reduced homology of the order complex of the open interval $(\hat{0}, v)$:*

$$b_{i,\mu}(I) = \tilde{H}_{i-1}(\hat{0}, v).$$

All other nontrivial multigraded Betti numbers are zero.

3. LATTICES SUPPORTING A PURE RESOLUTION

In this section, we prove our main theorem, characterizing the lattices supporting a pure resolution.

Definition 3.1. We say that Λ is *homologically monotonic* or *HM* if whenever $v < w \in \Lambda$ with $\tilde{H}_i(\hat{0}, v) \neq 0 \neq \tilde{H}_j(\hat{0}, w)$, then $i < j$.

For a nonnegative integer k , we say that Λ is *k -HM* if the above condition holds whenever $i \geq k$. (So the notion of “HM” is the same as “0-HM.”)

Remark 3.2. If a lattice Λ has a Cohen-Macaulay order complex, then it is HM. Recall Reisner’s criterion: For any x in such a lattice, $\tilde{H}_i(\hat{0}, x) = 0$ whenever i is less than the dimension of the order complex of $(\hat{0}, x)$. But if $x, y \in \Lambda$ with $x < y$, the dimension of $(\hat{0}, x)$ is strictly less than the dimension of $(\hat{0}, y)$. Thus, if we have $\tilde{H}_i(\hat{0}, x) \neq 0 \neq \tilde{H}_j(\hat{0}, y)$, it follows that i and j are the dimensions of the order complexes of $(\hat{0}, x)$ and $(\hat{0}, y)$, respectively, so $i < j$.

As shellable lattices are Cohen-Macaulay, this observation shows that many classes of lattices are HM, such as supersolvable lattices (which include distributive lattices) and geometric lattices.

Lemma 3.3. *Suppose that Λ is k -HM, and let $v \in \Lambda$ with $\tilde{H}_i(\hat{0}, v) \neq 0$ for some $i \geq k$. Then, for all other $j \geq k$, we have $\tilde{H}_j(\hat{0}, v) = 0$.*

Proof. Assume without loss of generality that $i < j$. Now let I be any ideal having lcm lattice Λ . (For example, I could be the Phan ideal of Λ , obtained by tagging each meet-irreducible element with a distinct variable, and tagging all other elements with 1, then

applying Construction 2.6.) Let

$$\dots \longrightarrow F_{j+1} \xrightarrow{\partial_{j+1}} F_j \longrightarrow \dots \longrightarrow F_i \longrightarrow \dots \longrightarrow F_0 \longrightarrow I \longrightarrow 0$$

be a minimal free resolution of I . If $\tilde{H}_j(\hat{0}, v) \neq 0$, then by Hochster's formula, F_{j+1} has a basis element e with multidegree $\tau^m(v)$. Then $\partial_{j+1}(e) \neq 0$, so we may write it as $\partial_{j+1}(e) = \sum m_f f$ for nonzero monomials m_f and basis elements f for F_j . Choose one such f ; we have $\text{mdeg}(m_f f) = \text{mdeg}(e) = \tau^m(v)$. It follows that $\text{mdeg}(f)$ divides $\tau^m(v)$ and that there exists $w \in \Lambda$ with $\tau^m(w) = \text{mdeg}(f)$, so $w < v$. But now we have $\tilde{H}_{j-1}(\hat{0}, w) \neq 0$, and $j-1 \geq i$, contradicting the assumption that Λ was k -HM. \square

Definition 3.4. Let (Λ, τ) be a tagged lattice. We say that (Λ, τ) is k -pure if whenever $\tilde{H}_k(\hat{0}, v) \neq 0 \neq \tilde{H}_k(\hat{0}, w)$ for two elements $v, w \in \Lambda$ then $\deg(\tau^m(v)) = \deg(\tau^m(w))$.

We also say that Λ is k -pure if there exists a tagging τ so that (Λ, τ) is k -pure. Finally, we say that (Λ, τ) (resp. Λ) is *pure* if it is k -pure for all $k \geq 0$.

In order to quantify how far a given tagged lattice (Λ, τ) is from being k -pure, it helps to define the statistic δ_k as follows.

Definition 3.5. For a tagged lattice (Λ, τ) , set

$$\delta_k(\Lambda, \tau) = \max\{\deg(\tau^m(v)) : \tilde{H}_k(\hat{0}, v) \neq 0\} - \min\{\deg(\tau^m(v)) : \tilde{H}_k(\hat{0}, v) \neq 0\}.$$

Note that (Λ, τ) is k -pure if and only if $\delta_k(\Lambda, \tau) = 0$.

Lemma 3.6. *Let (Λ, τ) be a tagged lattice that is not k -pure, and suppose Λ is k -HM. Then there exists a tagging τ_1 with the following two properties:*

- (1) $\delta_k(\Lambda, \tau_1) < \delta_k(\Lambda, \tau)$
- (2) If (Λ, τ) is ℓ -pure for some $\ell > k$, so is (Λ, τ_1) .

Proof. Define two subsets $S, T \subseteq \Lambda$ by

$$S = \{v \in \Lambda : \tilde{H}_k(\hat{0}, v) \neq 0 \text{ and } \deg(\tau^m(v)) \text{ is maximal}\},$$

$$T = \{v \in \Lambda : \tilde{H}_k(\hat{0}, v) \neq 0 \text{ and } \deg(\tau^m(v)) \text{ is not maximal}\}.$$

Because (Λ, τ) is not k -pure, both S and T are nonempty. We claim that no two elements $v, w \in S \cup T$ can be comparable (i.e., $S \cup T$ is an antichain of Λ). Indeed, if $v, w \in S \cup T$ were comparable (suppose $v < w$), we would have that both $\tilde{H}_k(\hat{0}, v)$ and $\tilde{H}_k(\hat{0}, w)$ are nonzero, but $k \not\leq k$, contradicting the assumption that Λ is k -HM.

Now write $S = \{v_1, v_2, \dots, v_t\}$, and let x_1, x_2, \dots, x_t be variables not appearing in the original tagging τ . Define τ_1 by $\tau_1(v) = \tau(v)$ if $v \notin S$ and $\tau_1(v_i) = \tau(v_i)x_i$ for $v_i \in S$. Now let $v \in S \cup T$. Since no two elements of $S \cup T$ are comparable, every element w of $S \cup T$ satisfies $v \not\leq w$, except for $w = v$. It follows that for $v \in T$ we have

$$\tau_1^m(v) = \tau^m(v)x_1x_2 \cdots x_t,$$

whereas for $v_i \in S$ we get

$$\tau_1^m(v_i) = \tau^m(v)x_1x_2 \cdots \widehat{x_i} \cdots x_t.$$

Thus, $\deg(\tau_1^m(v)) = \deg(\tau^m(v)) + t$ if $v \in T$, and $\deg(\tau_1^m(v)) = \deg(\tau^m(v)) + t - 1$ if $v \in S$, meaning $\delta_k(\Lambda, \tau_1) = \delta_k(\Lambda, \tau) - 1$. This proves (1).

Now we show (2). Choose $v, w \in \Lambda$ with $\tilde{H}_\ell(\hat{0}, v) \neq 0 \neq \tilde{H}_\ell(\hat{0}, w)$. Since (Λ, τ) is ℓ -pure, we have $\deg(\tau^m(v)) = \deg(\tau^m(w))$.

We claim that no $u \in S$ can satisfy $v \leq u$. Since $\tilde{H}_k(\hat{0}, u) \neq 0 \neq \tilde{H}_\ell(\hat{0}, v)$, we could not have $v < u$ since $k > \ell$ would contradict our assumption that Λ is k -HM. Similarly, by Lemma 3.3, we could not have $v = u$. Thus,

$$\deg(\tau_1^m(v)) = \deg(\tau^m(v)) + |\{u \in S : v \not\leq u\}| = \deg(\tau^m(v)) + |S|,$$

and the same holds for w : $\deg(\tau_1^m(w)) = \deg(\tau^m(w)) + |S|$. So we have $\deg(\tau_1^m(v)) = \deg(\tau_1^m(w))$, meaning (Λ, τ_1) remains ℓ -pure. \square

Lemma 3.7. *Let (Λ, τ) be a tagged lattice that is ℓ -pure, and suppose Λ is k -HM for some $k < \ell$. Then there is a tagging of Λ so that the associated tagged lattice is both k -pure and ℓ -pure.*

Proof. If (Λ, τ) is k -pure, we're done. Otherwise, we have $\delta_k(\Lambda, \tau) > 0$. We apply Lemma 3.6, obtaining a tagging τ_1 so that (Λ, τ_1) is ℓ -pure and $\delta_k(\Lambda, \tau_1) = \delta_k(\Lambda, \tau) - 1$. If $\delta_k(\Lambda, \tau_1) > 0$, we repeat the algorithm from Lemma 3.6 again, obtaining a tagging τ_2 so that (Λ, τ_2) is ℓ -pure and $\delta_k(\Lambda, \tau_2) = \delta_k(\Lambda, \tau_1) - 1$. Continuing, we eventually obtain a tagging $\sigma = \tau_j$ (actually, $j = \delta_k(\Lambda, \tau)$) so that (Λ, τ_j) is ℓ -pure and $\delta_k(\Lambda, \tau_j) = 0$, meaning (Λ, τ_j) is also k -pure. \square

Theorem 3.8. *Let Λ be an atomic lattice. Then there exists an ideal I with pure resolution such that $\text{lcm}(I) = \Lambda$ if and only if Λ is HM.*

Proof. First, suppose that Λ is not HM. Then there must be some $v, w \in \Lambda$ with $v < w$ and $\tilde{H}_i(\hat{0}, v) \neq 0 \neq \tilde{H}_j(\hat{0}, w)$ but $i \geq j$. If I is any ideal with $\text{lcm}(I) = \Lambda$, then v and w must correspond to monomials m_v and m_w with $\deg(m_v) < \deg(m_w)$. If $i = j$, then I does not have a pure resolution. If $i > j$, then $b_{i+1}(m_w) \neq 0$. It follows that $b_{j+1}(\mu) \neq 0$ for some μ dividing m_w . Since μ divides m_v , it follows that I does not have pure resolution.

Now suppose Λ is HM, and fix a tagging τ . Let i be the greatest integer for which (Λ, τ) is not i -pure. By Lemma 3.7, we can construct a tagging τ_1 so that (Λ, τ_1) is k -pure for all $k \geq i$. Now let j be the greatest integer for which (Λ, τ_1) is not pure (so that $j < i$). Again by Lemma 3.7, we can construct a tagging τ_2 such that (Λ, τ_2) is k -pure for all $k \geq j$. Continuing in this way, we eventually obtain a tagging τ_r so that (Λ, τ_r) is pure. Thus $I = I(\Lambda, \tau_r)$ is an ideal with pure resolution satisfying $\text{lcm}(I) = \Lambda$. \square

Example 3.9. If we apply the algorithm implicit in the proof of Theorem 3.8 to the running example of Section 2, the tagged lattice of Figure 2 becomes the new tagged lattice shown in Figure 4. The new variables in the tags are shown with capital letters.




FIGURE 4. The purification of the tagged lattice in Figure 2.

The corresponding monomial ideal is

$$(c^2dhijklm, acdefghkm, a^2b^2ijklm, abcfgijl),$$

which has a pure resolution in degrees 9, 13, and 16.

Observe that we could have done this with fewer variables by using pure powers for our new tags. Then the resulting monomial ideal is $(c^2dgh^3i^2, acdef^2gi^2, a^2b^2h^3i^2, abc f^2h^3)$. We could have used fewer variables still, if we were willing to reuse preexisting variables for our new tags, but the result is highly nonunique.

4. REMARKS AND QUESTIONS

Say that a monomial ideal is *purifiable* if there exists another ideal with pure resolution and the same lcm lattice. Then I is purifiable if and only if its lcm lattice is HM, and many important classes of monomial ideals are purifiable. It is natural from an algebraic perspective to seek out classes of ideals which are (or are not) purifiable.

Example 4.1. Recall that a monomial ideal is called *Scarf* if it is resolved by its Scarf complex; see [BPS] for the construction. Then, following the same reasoning as in Remark 3.2, Scarf ideals are purifiable: an element of the lcm lattice supports homology only if it appears in the supporting complex, and then only in the dimension of the corresponding cell. Since the “generic monomial ideals” defined in [BPS] are all Scarf, it follows that purifiability is very common.

A larger, if less well-studied, class of purifiable ideals is the *rigid* monomial ideals introduced by Clark and Mapes [CM]. These are the ideals whose resolutions have only scalar automorphisms; in particular, Scarf ideals are rigid. Clark and Mapes show that rigidity is equivalent to a stronger condition on the multigraded Betti numbers than being HM.

Definition 4.2 ([CM]). A monomial ideal I is *rigid* if it satisfies the following conditions.

- (R1) Every multigraded Betti number $b_{i,m}(I)$ is equal to either zero or one.

(R2) Multidegrees with the same nonzero Betti number are incomparable: If m divides m' , then for all i at least one of $b_{i,m}(I)$ and $b_{i,m'}(I)$ is equal to zero.

Applying Hochster's formula Theorem 2.11, it is clear that rigid ideals have HM lcm lattices and thus are purifiable. In fact, purifiability is equivalent to condition (R2):

Proposition 4.3. *A finite atomic lattice Λ is HM if and only if it satisfies the following condition: Suppose $v < w \in \Lambda$. Then for all i , at least one of $\tilde{H}_i(\hat{0}, v)$ and $\tilde{H}_i(\hat{0}, w)$ is zero.*

Proof. If Λ is HM, the given condition is immediate.

For the converse, the idea is the same as the proof of Lemma 3.3. Let I be any ideal with lcm lattice Λ , and let m_v and m_w be the corresponding multidegrees. Consider a minimal free resolution of I . The i^{th} free module has a generator in multidegree m_v if and only if $b_{i,m_v}(I) \neq 0$, if and only if $\tilde{H}_{i-1}(\hat{0}, m_v) \neq 0$. Now if Λ is not HM, there exist numbers $i \geq j$ and vertices $v < w \in \Lambda$ with $b_{i,m_v}(I)$ and $b_{j,m_w}(I)$ both nonzero. Choose a generator for the i^{th} free module with multidegree m_v , and apply the differential map. The result is an R -linear combination of generators for the $(i-1)^{\text{th}}$ free module, each with multidegree dividing m_v . Pick one such generator and repeat; after $(i-j)$ steps we will have found a generator for the j^{th} free module with multidegree strictly dividing m_w , violating the given condition. \square

In another direction, we can attempt to generalize the Scarf condition to a larger class of supporting complexes.

Question 4.4. Suppose that a minimal resolution of I is supported by a topological object X with nice structure. Are there conditions on X (e.g., being a regular cell complex) which allow us to conclude that I is purifiable?

Our intuition here is along the lines of Remark 3.2 and Example 4.1: each cell should correspond to a vertex in the lcm lattice, supporting homology only in its top dimension. The problem is that the face poset of X is not just a subset of the lcm lattice of I : faces can share multidegree, and incomparable faces can have comparable multidegrees.

We conclude with a remark about the role of \mathbb{k} .

Remark 4.5. Both purifiability of an ideal and homological monotonicity of a lattice are characteristic-dependent phenomena. Let $I = (abc, abf, ace, ade, adf, bcd, bde, bef, cdf, cef)$ be the Stanley-Reisner ideal of the minimal triangulation of the real projective plane, and let Λ be the lcm lattice of I . In characteristic other than two, I has a pure resolution, so Λ is HM. However, in characteristic two, I has multiple Betti numbers in multidegree $abcdef$. This corresponds to the open interval $(\hat{0}, \hat{1})$ having nonzero first and second homology in Λ , which means that Λ is not HM.

In fact, if Λ is any lattice with characteristic-dependent homology, it cannot be HM in the special characteristics: Looking at the resolution of any corresponding ideal, the consecutive cancellations that occur when passing from the special characteristic to characteristic zero correspond to a trivial complex in some (fixed) multidegree. The vertex of Λ with that multidegree consequently has two nonzero homologies in the special characteristic.

Acknowledgments: This work was partially supported by grants from the Simons Foundation (#199124 to Christopher Francisco and #202115 to Jeffrey Mermin).

REFERENCES

- [BCP] D. Bayer, H. Charalambous, and S. Popescu, Extremal Betti numbers and applications to monomial ideals. *J. Algebra* **221** (1999), no. 2, 497–512.
- [BPS] D. Bayer, I. Peeva, and B. Sturmfels, Monomial resolutions. *Math. Res. Lett.* **5** (1998), no. 1–2, 31–46.
- [CM] T. B. P. Clark and S. Mapes, Rigid monomial ideals. *J. Commut. Algebra* **6** (2014), no. 1, 33–52.
- [GPW] V. Gasharov, I. Peeva, and V. Welker, The lcm-lattice in monomial resolutions. *Math. Res. Lett.* **6** (1999), no. 5–6, 521–532.
- [H] M. Hochster, Cohen-Macaulay rings, combinatorics, and simplicial complexes. *Ring theory, II (Proc. Second Conf., Univ. Oklahoma, Norman, Okla., 1975)*, pp. 171–223. Lecture Notes in Pure and Appl. Math., Vol. 26, Dekker, New York, 1977.
- [M] S. Mapes, Finite atomic lattices and resolutions of monomial ideals. *J. Algebra* **379** (2013), 259–276.
- [Pe] I. Peeva, Graded Syzygies. *Algebra and Applications*, Volume 14. Springer, London (2011).
- [Ph] J. P. Phan, Order properties of monomial ideals and their free resolutions. Thesis (Ph.D.), Columbia University. 2006. 54 pp.
- [V] M. Velasco, Minimal free resolutions that are not supported by a CW-complex. *J. Algebra* **319** (2008), no. 1, 102–114.

DEPARTMENT OF MATHEMATICS, OKLAHOMA STATE UNIVERSITY, 401 MATHEMATICAL SCIENCES,
STILLWATER, OK 74078

E-mail address: `chris.francisco@okstate.edu`

URL: `http://math.okstate.edu/people/chris/`

DEPARTMENT OF MATHEMATICS, OKLAHOMA STATE UNIVERSITY, 401 MATHEMATICAL SCIENCES,
STILLWATER, OK 74078

E-mail address: `mermin@math.okstate.edu`

URL: `http://math.okstate.edu/people/mermin/`

DEPARTMENT OF MATHEMATICS, OKLAHOMA STATE UNIVERSITY, 401 MATHEMATICAL SCIENCES,
STILLWATER, OK 74078

E-mail address: `jay.schweig@okstate.edu`

URL: `http://math.okstate.edu/people/jayjs/`